


# Unlike *Drosophila elav*, the *C. elegans elav* orthologue *exc-7* is not panneuronally expressed

Kenneth Pham<sup>1</sup> and Oliver Hobert<sup>1</sup>

<sup>1</sup> Columbia University, Department of Biological Sciences, HHMI, New York, NY

To whom correspondence should be addressed: or38@columbia.edu



expression. We cloned *gfp* from the *che-1(ot856[che-1::gfp])* allele (Leyva-Díaz and Hobert, 2019), inserted it into the pMiniT 2.0 vector (NEB), and used that resulting plasmid for subsequent cloning of the *gfp* tag.

Embryonic expression of *exc-7* was first observed at the bean stage. By reverse lineaging with use of SIMI-Biocell software (Schnabel *et al.*, 1997), we confirm the identity of one of the expressing cells at this stage as the excretory canal cell (Fig. 1B, arrow). In L1 animals, broad expression in the head, ventral nerve cord (VNC), and tail was observed (Fig. 1C). In young adults, expression is notably observed in vulva cells (Fig. 1D’). In the nervous system specifically, expression is observed in many neurons throughout the body (Fig. 1D’-D’’), but unlike *Drosophila* *Elav*, *exc-7::gfp* it is not panneuronally expressed. We used the NeuroPAL transgene (<https://www.biorxiv.org/content/10.1101/676312v1>) to individually identify each neuron in which *exc-7* is expressed in the young adult worm. Sites of expression are listed in Fig. 1F and some examples of neuronal expression are shown in Fig. 1E’. Expression in all neurons is at least several fold more intense than UPN::NLS::TagRFP-T signal from NeuroPAL. We confirmed previously reported expression in cholinergic VNC MNs, but absence of GABAergic VNC MNs (Fig. 1E’), consistent with previous reports (Fujita *et al.*, 1999; Loria *et al.*, 2003) and consistent with *exc-7* functioning in cholinergic, but not GABAergic neurons to control alternative splicing (Norris *et al.*, 2014). *exc-7::gfp* is also expressed in some non-neuronal cell types, including muscle and hypodermis, but not in the gut (Fig. 1D’-D’’). A previous report showed that *exc-7* is only transiently and weakly expressed in the excretory cell, which, based on *exc-7*'s excretory mutant phenotype, has puzzled researchers (Fujita *et al.*, 2003). We find that the *gfp* tagged *exc-7* locus is strongly and continuously expressed in the excretory canal cell (Fig. 1B-D’, arrow). We conclude that unlike its fly orthologue *elav*, *exc-7* is not a panneuronally expressed gene.

## Reagents

OH16020: *exc-7(ot970[exc-7::gfp::3xflag])*

The strain is available through the CGC.

**Acknowledgments:** We thank Chi Chen for expert DNA injection, Eduardo Leyva-Díaz for advice with CRISPR genome editing, and Neda Masoudi for help with embryonic lineaging.

## References

- Berger, C., Renner, S., Luer, K., and Technau, G.M. (2007). The commonly used marker ELAV is transiently expressed in neuroblasts and glial cells in the *Drosophila* embryonic CNS. *Dev Dyn* 236, 3562-3568. PMID: 17994541.
- Campos, A.R., Rosen, D.R., Robinow, S.N., and White, K. (1987). Molecular analysis of the locus *elav* in *Drosophila melanogaster*: a gene whose embryonic expression is neural specific. *EMBO J* 6, 425-431. PMID: 3107982.
- Dokshin, G.A., Ghanta, K.S., Piscopo, K.M., and Mello, C.C. (2018). Robust Genome Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in *Caenorhabditis elegans*. *Genetics* 210, 781-787. PMID: 30213854.
- Fujita, M., Hawkinson, D., King, K.V., Hall, D.H., Sakamoto, H., and Buechner, M. (2003). The role of the ELAV homologue EXC-7 in the development of the *Caenorhabditis elegans* excretory canal. *Dev Biol* 256, 290-301. PMID: 12679103.
- Fujita, M., Kawano, T., Ohta, A., and Sakamoto, H. (1999). Neuronal expression of a *Caenorhabditis elegans* *elav*-like gene and the effect of its ectopic expression. *Biochem Biophys Res Commun* 260, 646-652. PMID: 10403820.
- Leyva-Díaz, E., and Hobert, O. (2019). Transcription factor autoregulation is required for acquisition and maintenance of neuronal identity. *Development* 146. PMID: 31227642.
- Loria, P.M., Duke, A., Rand, J.B., and Hobert, O. (2003). Two neuronal, nuclear-localized RNA binding proteins involved in synaptic transmission. *Curr Biol* 13, 1317-1323. PMID: 12906792.
- Luo, L., Liao, Y.J., Jan, L.Y., and Jan, Y.N. (1994). Distinct morphogenetic functions of similar small GTPases: *Drosophila* *Drac1* is involved in axonal outgrowth and myoblast fusion. *Genes Dev* 8, 1787-1802. PMID: 7958857.
- Norris, A.D., Gao, S., Norris, M.L., Ray, D., Ramani, A.K., Fraser, A.G., Morris, Q., Hughes, T.R., Zhen, M., and Calarco, J.A. (2014). A pair of RNA-binding proteins controls networks of splicing events contributing to specialization of neural cell types. *Mol Cell* 54, 946-959. PMID: 24910101.
- O'Neill, E.M., Rebay, I., Tjian, R., and Rubin, G.M. (1994). The activities of two Ets-related transcription factors required for *Drosophila* eye development are modulated by the Ras/MAPK pathway. *Cell* 78, 137-147. PMID: 8033205.
- Osterwalder, T., Yoon, K.S., White, B.H., and Keshishian, H. (2001). A conditional tissue-specific transgene expression system using inducible GAL4. *Proc Natl Acad Sci U S A* 98, 12596-12601. PMID: 11675495.
- Robinow, S., and White, K. (1988). The locus *elav* of *Drosophila melanogaster* is expressed in neurons at all developmental stages. *Dev Biol* 126, 294-303. PMID: 3127258.

Robinow, S., and White, K. (1991). Characterization and spatial distribution of the ELAV protein during *Drosophila melanogaster* development. *J Neurobiol* 22, 443-461. PMID: 1716300.

Samson, M.L. (2008). Rapid functional diversification in the structurally conserved ELAV family of neuronal RNA binding proteins. *BMC Genomics* 9, 392. PMID: 18715504.

Schnabel, R., Hutter, H., Moerman, D., and Schnabel, H. (1997). Assessing normal embryogenesis in *Caenorhabditis elegans* using a 4D microscope: variability of development and regional specification. *Dev Biol* 184, 234-265. PMID: 9133433.

Stefanakakis, N., Carrera, I., and Hobert, O. (2015). Regulatory Logic of Pan-Neuronal Gene Expression in *C. elegans*. *Neuron* 87, 733-750. PMID: 26291158.

**Funding:** This work was supported by the HHMI.

**Author Contributions:** Kenneth Pham: Investigation, Methodology, Visualization, Writing - review and editing. Oliver Hobert: Funding acquisition, Supervision, Writing - original draft.

**Reviewed By:** Matthew Buechner

**History: Received** October 19, 2019 **Accepted** October 30, 2019 **Published** October 30, 2019

**Copyright:** © 2019 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Pham, K; Hobert, O (2019). Unlike *Drosophila elav*, the *C. elegans* *elav* orthologue *exc-7* is not panneuronally expressed. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000189>