

Caenorhabditis Intervention Testing Program: the creatine analog β -guanidinopropionic acid does not extend lifespan in nematodes

Anna L. Coleman-Hulbert¹, Erik Johnson¹, Christine A. Sedore¹, Stephen A. Banse¹, Max Guo², Monica Driscoll³, Gordon J. Lithgow⁴ and Patrick C. Phillips^{1§}


¹Institute of Ecology and Evolution, University of Oregon, Eugene, Oregon 97403, USA

²Division of Aging Biology, National Institute on Aging, Bethesda, Maryland 20892, USA

³Department of Molecular Biology and Biochemistry, Rutgers University, Piscataway, New Jersey 08854, USA

⁴The Buck Institute for Research on Aging, Novato, California 94945, USA

[§]To whom correspondence should be addressed: pphil@uoregon.edu



Longevity under adult β -guanidinopropionic acid exposure: Survival curves for *C. elegans* strain N2, *C. briggsae* strain AF16, and *C. tropicalis* strain JU1373 exposed to β -guanidinopropionic acid at various concentrations starting on the first day of adulthood. Only the lifespan of AF16 at 300 mM (mean=26.2 days, $p=0.0123$) differed significantly from the control (mean=31.5 days for AF16; control means for N2=18.8 days and JU1373=25.1 days; statistical comparisons were made with a Cox proportional hazards mixed-model using the `coxme` v.2.2-5 package in R (Therneau 2012; Lucanic *et al.* 2017)).

Description

The *Caenorhabditis* Intervention Testing Program (CITP) is a multi-institutional, National Institutes of Aging (NIA)-funded consortium charged with identifying chemical compounds that robustly extend lifespan in a genetically diverse panel of *Caenorhabditis* strains. Compounds are prioritized for screening if they are highly ranked via computational prediction for lifespan or healthspan effects (Coleman-Hulbert *et al.* 2019), if they are predicted to engage known lifespan regulating pathways, or if they have previously been reported as extending lifespan or healthspan in model systems (Lucanic *et al.* 2017). β -guanidinopropionic acid (β -GPA) is a creatine analog (Shields and Whitehair 1973), commonly used as a dietary supplement, and has been shown to extend lifespan in *Drosophila* under stress via 5' AMP-activated protein kinase (AMPK) activity (Yang *et al.* 2015). The AMPK pathway is conserved in nematodes and humans (Apfeld *et al.* 2004) and is involved in multiple pathways affecting stress response and metabolism (Wang *et al.* 2012).

We assayed lifespan in response to β -GPA exposure in three *Caenorhabditis* species using our previously published workflow (Lucanic *et al.* 2017). In brief, worms were age-synchronized by timed egg-lays on standard 60 mm diameter Nematode Growth Media (NGM) plates and transferred at a density of 50 individuals per 35 mm treated plate in triplicate when they reached adulthood (for control plates, there were six replicates of 50 animals each). β -GPA (Sigma-Aldrich) was dissolved in water and diluted appropriately such that addition of 125 μ l of solution to 35 mm diameter plates containing NGM with lawns of *E. coli* OP50-1 and 51 μ m FUDR would generate the following final β -GPA concentrations: 0.1 μ M, 1 μ M, 10 μ M, 50 μ M, 100 μ M, 500 μ M, 1 mM, 10 mM and 300 mM. Worms were maintained at 20 °C and moved to fresh plates on the first, second, and fourth (*C. tropicalis*) or fifth (*C. elegans* and *C. briggsae*) day of adulthood, then once weekly afterward. Due to our previous experience with compound interventions that potentially alter bacterial viability through transient pH changes upon plate treatment (Banse *et al.* 2019), we investigated the effects of β -GPA on our assay plates. At the above listed concentrations, β -GPA-treated plates had a pH of 6.5 and the bacterial lawns survived treatment when tested by replica plating. Thrice weekly, we observed animals for spontaneous movement or movement after gentle perturbation with a 0.2 mm diameter platinum wire. Death was scored as a lack of movement.

Our results indicate that β -GPA does not extend lifespan in the three nematode species at the concentrations tested here; in fact, in only one instance was an effect detected and, in that case, the compound reduced lifespan (Fig. 1). This conclusion is based on one biological replicate per concentration and, as such, could be considered preliminary. While interventions may be ineffective due to a range of causes, including permeability barriers, compound stability *in vivo*, and metabolism by the bacterial food source, we believe that the lack of response in this study was due to a lack of physiological relevance

in *C. elegans*. In *Drosophila* and mammalian models, β -GPA reduces the level of intracellular phosphocreatine that can be used by creatine kinases to regenerate ATP (Oudman *et al.* 2013; Yang *et al.* 2015) resulting in a decreased cellular ATP/AMP ratio, which activates AMPK and ultimately increases lifespan and stress-resistance. Our *a priori* expectation for lifespan extension in *C. elegans* was built on the observations that: (1) creatine is reported as detectable in *C. elegans* (Atherton *et al.* 2008; Jones *et al.* 2012; Wan *et al.* 2017); (2) *C. elegans* has a creatine-like kinase, ARGK-1, whose activity modulates AMPK signaling (McQuary *et al.* 2016); (3) modulation of AMPK (Apfeld *et al.* 2004; Greer *et al.* 2007) and ARGK-1 (McQuary *et al.* 2016) in *C. elegans* can affect lifespan and stress resistance; and (4) regulation of lifespan by insulin signaling is partially dependent on AMPK signaling in *C. elegans* (Tullet *et al.* 2014). As such, we were surprised to find no changes in lifespan upon treatment with a creatine analog. One possible explanation is that, despite the similarity between ARGK-1 and mammalian creatine kinase, the enzymes' substrates differ. Biochemical analysis suggests that ARGK-1 uses arginine instead of creatine as a substrate to recharge ADP (Fraga *et al.* 2015). Additionally, it has been postulated that the biochemical characterization of *C. elegans* metabolites may have misidentified creatine, and that creatine is not relevant to *C. elegans* physiology (Witting *et al.* 2018). Given these caveats, it may not be surprising that β -GPA does not alter *Caenorhabditis* lifespan.

Acknowledgments: We acknowledge all of the members of the Lithgow, Driscoll and Phillips labs for helpful discussions. We thank the CITP Advisory Committee and Ronald Kohanski (National Institute on Aging) for extensive discussion. Strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

References

- Apfeld, J., O'Connor, G., McDonagh, T., DiStefano, P.S., and Curtis, R. (2004). The AMP-activated protein kinase AAK-2 links energy levels and insulin-like signals to lifespan in *C. elegans*. *Genes Dev* 18, 3004–3009. PMID: 15574588.
- Atherton, H.J., Jones, O.A.H., Malik, S., Miska, E.A., and Griffin, J.L. (2008). A comparative metabolomic study of NHR-49 in *Caenorhabditis elegans* and PPAR- α in the mouse. *FEBS Lett* 582, 1661–1666. PMID: 18435929.
- Banse S.A., Lucanic, M., Sedore, C.A., Coleman-Hulbert, A.L., Plummer, W.T., Chen, E., Kish, J.L., Hall, D., Onken, B., Presley, M.P., Jones, E.G., Blue, B.W., Garrett, T., Abbott, M., Xue, J., Guo, S., Johnson, E., Foulger, A.C., Chamoli, M., Falkowski, R., Melentijevic, I., Harinath, G., Huynh, P., Patel, S., Edgar, D., Jarrett, C.M., Guo, M., Kapahi, P., Lithgow, G.J., Driscoll, M., and Phillips, P.C. (2019). Automated lifespan determination across *Caenorhabditis* strains and species reveals assay-specific effects of chemical interventions. *GeroScience*. PMID: 31820364.
- Coleman-Hulbert, A.L., Johnson, E., Sedore, C.A., Banse, S.A., Guo, M., Driscoll, M., Lithgow, G.J., and Phillips, P.C. (2019). *Caenorhabditis* Intervention Testing Program: the tyrosine kinase inhibitor imatinib mesylate does not extend lifespan in nematodes. *microPublication Biol*. DOI: 10.17912/micropub.biology.000131
- Fraga, D., Aryal, M., Hall, J.E., Rae, E., and Snider, M. (2015). Characterization of the arginine kinase isoforms in *Caenorhabditis elegans*. *Comp Biochem Physiol B Biochem Mol Biol* 187, 85–101. PMID: 25981702.
- Greer, E.L., Dowlatshahi, D., Banko, M.R., Villen J., Hoang, K., Blanchard, D., Gygi, S.P., and Brunet, A. (2007). An AMPK-FOXO pathway mediates longevity induced by a novel method of dietary restriction in *C. elegans*. *Curr Biol* 17, 1646–1656. PMID: 17900900.
- Jones, O.A.H., Swain, S.C., Svendsen, C., Griffin, J.L., Sturzenbaum, S.R., and Spurgeon, D.J. (2012). Potential new method of mixture effects testing using metabolomics and *Caenorhabditis elegans*. *J Proteome Res* 11, 1446–1453. PMID: 22175231.
- Lucanic, M., Plummer, W.T., Chen, E., Harke, J., Foulger, A.C., Onken, B., Coleman-Hulbert, A.L., Dumas, K.J., Guo, S., Johnson, E., Bhaumik, D., Xue, J., Crist, A.B., Presley, M.P., Harinath, G., Sedore, C.A., Chamoli, M., Kamat, S., Chen, M.K., Angeli, S., Chang, C., Willis, J.H., Edgar, D., Royal, M.A., Chao E.A., Patel, S., Garrett, T., Ibanez-Ventoso, C., Hope, J., Kish, J.L., Guo, M., Lithgow, G.J., Driscoll, M., and Phillips, P.C. (2017). Impact of genetic background and experimental reproducibility on identifying chemical compounds with robust longevity effects. *Nat Commun* 8, 14256. PMID: 28220799.
- McQuary, P.R., Liao, C.-Y., Chang, J.T., Kumsta, C., She, X., Davis, A., Chu, C.-C., Gelino, S., Gomez-Amaro, R.L., Petrascheck, M., Brill, L.M., Ladiges, W.C., Kennedy, B.K., and Hansen, M. (2016). *C. elegans* S6K Mutants Require a Creatine-Kinase-like Effector for Lifespan Extension. *Cell Rep* 14, 2059–2067. PMID: 26923601.
- Oudman, I., Clark, J.F., and Brewster, L.M. (2013). The effect of the creatine analogue beta-guanidinopropionic acid on energy metabolism: a systematic review. *PLoS One* 8, e52879. PMID: 23326362.
- Shields, R.P., and Whitehair, C.K. (1973). Muscle creatine: in vivo depletion by feeding beta-guanidinopropionic acid. *Can J Biochem* 51, 1046–1049. PMID: 4725354.

Therneau, T. (2012). coxme: Mixed Effects Cox Models R Foundation for Statistical Computing R package version 2, 2–5.

Tullet, J.M.A., Araiz, C., Sanders, M.J., Au, C., Benedetto, A., Papatheodorou, I., Clark, E., Schmeisser, K., Jones, D., Schuster, E.F., Thornton, J.M., and Gems, D. (2014). DAF-16/FoxO directly regulates an atypical AMP-activated protein kinase gamma isoform to mediate the effects of insulin/IGF-1 signaling on aging in *Caenorhabditis elegans*. PLoS Genet 10, e1004109. PMID: 24516399.

Wan, Q.-L., Shi, X., Liu, J., Ding, A.-J., Pu, Y.-Z., Li, Z., Wu, G.-S., and Luo, H.-R. (2017). Metabolomic signature associated with reproduction-regulated aging in *Caenorhabditis elegans*. Aging (Albany, NY) 9, 447–474. PMID: 28177875.

Wang, S., Song, P., and Zou, M.-H. (2012). AMP-activated protein kinase, stress responses and cardiovascular diseases. Clin Sci (Lond) 122, 555–573. PMID: 22390198.

Witting, M., Hastings, J., Rodriguez, N., Joshi, C.J., Hattwell, J.P.N., Ebert, P.R., van Weeghel, M., Gao, A.W., Wakelam, M.J.O., Houtkooper, R.H., Mains, A., Le Novère, N., Sadykoff, S., Schroeder, F., Lewis, N.E., Schirra, H.-J., Kaleta, C., and Casanueva, O. (2018). Modeling Meets Metabolomics-The WormJam Consensus Model as Basis for Metabolic Studies in the Model Organism *Caenorhabditis elegans*. Front Mol Biosci 5, 96. PMID: 30488036.

Yang, S., Long, L.H., Li, D., Zhang, J.K., Jin, S., Wang, F., and Chen, J.G. (2015). β -Guanidinopropionic acid extends the lifespan of *Drosophila melanogaster* via an AMP-activated protein kinase-dependent increase in autophagy. Aging Cell 14, 1024–1033. PMID: 26120775.

Funding: This work was supported by National Institutes of Health grants U01 AG045864 to MD, U01 AG045844 to GJL, and U01 AG045829 and U24 AG056052 to PCP.

Author Contributions: Anna L. Coleman-Hulbert: Writing - original draft, Writing - review and editing, Resources, Project administration, Investigation. Erik Johnson: Investigation, Writing - review and editing. Christine A. Sedore: Investigation, Formal analysis, Writing - review and editing, Visualization. Stephen A. Banse: Methodology, Writing - original draft, Writing - review and editing. Max Guo: Conceptualization, Methodology, Writing - review and editing. Monica Driscoll: Funding acquisition, Methodology, Writing - review and editing. Gordon J. Lithgow: Funding acquisition, Methodology, Writing - review and editing. Patrick C. Phillips: Funding acquisition, Methodology, Supervision, Writing - review and editing.

Reviewed By: Anonymous

History: Received December 20, 2019 **Accepted** December 29, 2019 **Published** January 2, 2020

Copyright: © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Coleman-Hulbert, AL; Johnson, E; Sedore, CA; Banse, SA; Guo, M; Driscoll, M; Lithgow, GJ; Phillips, PC (2020). *Caenorhabditis* Intervention Testing Program: the creatine analog β -guanidinopropionic acid does not extend lifespan in nematodes. microPublication Biology. <https://doi.org/10.17912/micropub.biology.000207>