


# A role for *Arabidopsis* myosins in sugar-induced hypocotyl elongation

Damilola Olatunji<sup>1</sup> and Dior R Kelley<sup>1§</sup>

<sup>1</sup>Iowa State University

<sup>§</sup>To whom correspondence should be addressed: dkelley@iastate.edu

## Abstract



**Figure 1:** (A) Schematic gene structure of *ATM1* (AT3G19960) indicating the T-DNA insertions assayed in this study: SAIL\_405\_B08 (*atm1-1*), SALK\_127305 (*atm1-2*), GABI\_209H11 (*atm1-3*), and SALK\_022962 (*atm1-4*). Black boxes and lines depict exons and introns respectively. Scale bar is 100 bases. (B) *ATM1* transcript levels in different alleles (*atm1-1*, *atm1-2*, *atm1-3*, and *atm1-4*) compared to Columbia Ecotype (Col-0) assayed by quantitative PCR (qPCR). (C) Average hypocotyl length of seedlings grown on 0.5X Murashige and Skoog (MS) medium with or without 15 mM sucrose in the dark for 5 days. A total of 30-40 seedlings were evaluated and the assay was replicated three times (panel shows the results from one experiment). Error bars show standard error. P values are calculated using one- or two-way ANOVA followed by Tukey multiple comparisons; ns: not significant. (D) Hypocotyl phenotypes after 5 days of incubation in continuous darkness. Scale bar: 5mm

## Description

The ability of developing seedlings to respond and adapt to diverse environmental conditions including light is critical for their emergence and establishment (Benvenuti *et al.*, 2001; Forcella *et al.*, 2000; Salter *et al.*, 2003; Yu and Huang, 2017). Cell expansion within the hypocotyl optimizes light and energy capture by the cotyledons, and enables the transition to autotrophic status (Botterweg-Paredes *et al.*, 2020; Dowson-Day and Millar, 1999; Oh *et al.*, 2013). Hypocotyl elongation is regulated by multiple factors including temperature, phytohormones, circadian clock and light (Dowson-Day and Millar, 1999; Ma *et al.*, 2016; Procko *et al.*, 2014; Reed *et al.*, 2018; Yu and Huang, 2017). Endogenous and exogenous sugars are also important regulators of hypocotyl cell expansion (Lilley *et al.*, 2012; Liu *et al.*, 2011; Pfeiffer and Kutschera, 1995; Simon *et al.*, 2018b; Singh *et al.*, 2017; Zhang *et al.*, 2010, 2015). Under constant darkness, the interaction between plant hormones such as brassinosteroid and gibberellin and sugar signalling is proposed to stimulate increase in hypocotyl

length (Simon *et al.*, 2018b; Zhang *et al.*, 2010, 2015). Hypocotyl phenotypes are important tools in plant biology as they have been used to screen for mutants with altered responses to light and sugar signalling and some of the identified genes have revolutionized plant research (Nakano, 2019). In multicellular organisms, actomyosin-dependent transport constitutes an essential component of the cellular structure and dynamics (Duan and Tominaga, 2018; Kurth *et al.*, 2017; Peremyslov *et al.*, 2010). The Arabidopsis genome encodes 17 myosins classified into two distantly-related groups comprising 4 class VIII myosins and 13 class XI plant myosins (Haraguchi *et al.*, 2019; Reddy and Day, 2001; Ryan and Nebenführ, 2018). The plant-specific-myosin XI has been implicated in diverse developmental processes (Cai *et al.*, 2014), but the contribution of members of class VIII myosins including myosin1/ATM1 to plant development stills remains elusive.

Here, we analyzed the role of plant myosins in sucrose-induced hypocotyl elongation under continuous darkness conditions. To start the molecular characterization of *ATM1* (AT3G19960), we analyzed four T-DNA insertional alleles (Fig. 1A), and mutations were confirmed via PCR-based genotyping. We also performed RT-qPCR to quantify *ATM1* transcripts in the four alleles, of which *atm1-1* (SAIL\_405\_BO8 previously described by Haraguchi *et al.*, 2014) showed reduced *ATM1* expression among these alleles compared to the wild type (Fig. 1B). Notably, *atm1-3* exhibits increased mRNA levels and may be a overexpression allele. Since *atm1-1* had the lowest *ATM1* transcript level, and was previously described as a null (Haraguchi *et al.*, 2014) we decided to use this allele for further phenotypic analysis. To understand the contribution of plant myosins in sugar-regulated hypocotyl cell expansion we performed growth assays using Col-0 (WT: wild-type), *atm1-1*, a class XI myosin single gene mutant (*xik*) and the triple knockout line (*3KO*; *xi1 xi2 xik*). This triple mutant was previously generated using SALK insertion lines in XI-K (SALK\_067972; At5g20490), XI-1 (SALK\_019031; At1g17580) and XI-2 (SALK\_055785; At5g43900) (Prokhnevsky *et al.*, 2008; Valera *et al.*, 2010). All genotypes were grown on 0.5X Murashige and Skoog (MS) medium without or with 15 mM sucrose under constant darkness for 5 days. Under the sucrose-free condition (control), all the three myosin mutants showed reduced hypocotyl length compared to the WT plants (Fig. 1C and D), suggesting the role of myosins in cell elongation. Specifically, the *3KO* mutant was severely impaired in hypocotyl cell expansion in the absence of sucrose when compared to the single mutants, hinting that this gene family may redundantly regulate plant development. In the presence of sucrose under constant darkness, the short hypocotyl phenotype was not fully rescued when compared to the WT plants except for *atm1-1*.

Altogether our findings suggest that plant-specific myosins may be involved in sugar-regulated hypocotyl elongation under continuous darkness. These myosins have diverse subcellular localization patterns (Haraguchi *et al.*, 2014; Peremyslov *et al.*, 2012, 2008) and thus may represent a new downstream signaling output following sucrose signals within the developing shoot. Sucrose-induced hypocotyl elongation has been linked to brassinosteroid, gibberellins, phytochrome interacting factors (PIFs) and SnRK1 (Sucrose Nonfermenting Kinase 1) signaling (Laxmi *et al.*, 2004; Liu *et al.*, 2011; Simon *et al.*, 2018a; Zhang *et al.*, 2015, 2010; Zhang and He, 2015). Other proteins that are important for this process include TANG1 (Zheng *et al.*, 2015), a WD40 protein AtGHS40 (Hsiao *et al.*, 2016), a plant-specific protein SR45 (Carvalho *et al.*, 2010), and HIGH SUGAR RESPONSE 8 (HSR8) (Li *et al.*, 2007). Further studies will be needed to determine how these molecular factors work in concert or parallel to regulate skotomorphogenesis.

## Methods

[Request a detailed protocol](#)

### Plant materials

Arabidopsis thaliana plants used in this study were Columbia (Col-0) ecotype: SAIL\_405\_B08 (*atm1-1*), SALK\_127305C (*atm1-2*), GABI\_219H11(*atm1-3*) and SALK\_022962C (*atm1-4*) and previously described (*xik*) and *3KO* triple knockout line (*xi1 xi2 xik*) (Peremyslov *et al.*, 2012, 2010, 2008). Seeds were surfaced sterilized using 50% bleach and 0.01% Triton X-100 for 10 min and then washed five times with sterile water. Seeds were then imbibed in sterile water for 2 days at 4 °C and then transferred to 0.5X Murashige and Skoog (MS) medium plates supplemented without/with 15mM sucrose. For hypocotyl elongation, plates were incubated in the light for 6 hours prior to incubation in the dark for 5 days at 22 °C. PCR based genotyping of the mutants was performed with primers listed in Table 1 and 2X DreamTaq polymerase master mix (Thermo Fisher Scientific).

### Real time quantitative PCR (RT-qPCR)

Plant materials were harvested from 7-day old seedlings in three biological replicates. RNA samples were extracted with ZYMO RESEARCH Direct-zol RNA Miniprep Plus Kit and quality was spectrophotometrically measured with the Nanodrop. cDNA synthesis was performed with SuperScript IV Reverse Transcriptase kit according to manufacturer's instruction. The samples were run on *BIO RAD CFX Connect* Real-Time PCR Detection System with the following components per reaction of 20μL volume: 10μL iTaq Universal SYBR Green Supermix, 0.6 μL Forward Primer F (300nM), 0.6 μL Reverse Primer F (300nM) and 3μL cDNA (5ng/ul). No cDNA samples (water) were included as negative control. Cycling conditions were 5 min at 95 °C, followed by cycles of 15 s at 95 °C, 30 s at 60 °C and 30 s at 72 °C. Data acquisition was done at the end of every cycle. The samples were prepared in three biological and two technical

repeats. The Comparative CT Method ( $\Delta\Delta$  CT Method) was used for the analysis of Ct values whereby the amount of target, normalized to an endogenous reference and relative to a calibrator and is given by  $2^{-\Delta\Delta CT}$ .

### Hypocotyl length and statistical analysis

Five-day old dark-grown seedlings were imaged using an Epson Perfection V600 Photo Scanner. Hypocotyl length was measured with *ImageJ*. For each experimental treatment, 30-40 seedlings were measured for each genotype. The assays were repeated three independent times with similar results. GraphPad Prism (GraphPad Software, version 8.4.2) was used for statistical analysis. To compare treatments effects on the mean value of wild type and mutants, either one- or two-way ANOVA was performed using Tukey's test for multiple comparison.

### Gene diagram annotation

The diagram of *ATM1* (*AT3G19960*) was created using the Exon-Intron Graphic Maker (<http://wormweb.org/exonintron>) using current gene model sequence from TAIR ([www.arabidopsis.org](http://www.arabidopsis.org)). The position of all T-DNA alleles examined was obtained from the SALK SIGnAL T-DNA express Arabidopsis Gene Mapping tool (<http://signal.salk.edu/cgi-bin/tdnaexpress>).

## Reagents

**Table 1:** Primers used in this study.

Primers	5' > 3' Sequence	Purpose
SAIL_405_B08 LP	TTCGTGTGAACGTTGATTCTG	Genotyping
SAIL_405_B08 RP	TCCAGCTTGAATAGATGACGG	
SALK_127305_LP	TCCTCAAGCATCACCGTTAAC	
SALK_127305_RP	GCAGAGAGCTCAAGTGTGG	
GABI_219H11_LP	TAAGAGCGAGACAGAGAACCG	
GABI_219H11_RP	TCGTGGTTGGTTGGTTAGAAG	
SALK_022962_LP	GGGAAACAGAGAGAAATTGG	
SALK_022962_RP	TTTGCTTTGGCATTAAACCAAC	
Sail-LB2	GCTTCCTATTATATCTTCCCAAATTACCAATACA	
LBb1.3	ATTTTGCCGATTCGGAAC	
GABI-8474	ATAATAACGCTGCGGACATCTACATTTT	
ATM1_qPCR_Fw	CAGACAGAGAACTGAGGAGGC	RT-qPCR
ATM1_qPCR_Rv	CATCGAACCCTGCTCTCTTCG	
AT1G13320_Fw	GCGGTTGTGGAGAACATGATACG	
AT1G13320_Rv	GAACCAAACACAATTCGTTGCTG	

**Acknowledgments:** We wish to thank Valerian Dolja (Oregon State University) for sharing seed stocks of the previously described (xik) and 3KO triple knockout lines (xi1 xi2 xik) (Peremyslov et al., 2008). All other seed stocks were obtained from the ABRC (Arabidopsis Biological Resource Center at Ohio State University).

## References

- Benvenuti, S., Macchia, M., Miele, S., 2001. Light, temperature and burial depth effects on *Rumex obtusifolius* seed germination and emergence. *Weed Res.* 41, 177–186. DOI: <https://doi.org/10.1046/j.1365-3180.2001.00230.x>
- Botterweg-Paredes, E., Blaakmeer, A., Hong, S.-Y., Sun, B., Mineri, L., Kruusvee, V., Xie, Y., Straub, D., Ménard, D., Pesquet, E., Wenkel, S., 2020. Light affects tissue patterning of the hypocotyl in the shade-avoidance response. *PLOS Genet.* 16, e1008678. PMID: 32203519.
- Cai, C., Henty-Ridilla, J.L., Szymanski, D.B., Staiger, C.J., 2014. Arabidopsis Myosin XI: A Motor Rules the Tracks. *Plant Physiol.* 166, 1359 LP – 1370. PMID: 25237128.
- Carvalho, R.F., Carvalho, S.D., Duque, P., 2010. The plant-specific SR45 protein negatively regulates glucose and ABA signaling during early seedling development in Arabidopsis. *Plant Physiol.* 154, 772–783. PMID: 20699397.

- Dowson-Day, M.J., Millar, A.J., 1999. Circadian dysfunction causes aberrant hypocotyl elongation patterns in *Arabidopsis*. *Plant J.* 17, 63–71. PMID: 10069068.
- Duan, Z., Tominaga, M., 2018. Actin-myosin XI: an intracellular control network in plants. *Biochem. Biophys. Res. Commun.* 506, 403–408. PMID: 29307817.
- Forcella, F., Benech Arnold, R.L., Sanchez, R., Ghersa, C.M., 2000. Modeling seedling emergence. *F. Crop. Res.* 67, 123–139. DOI: [https://doi.org/10.1016/S0378-4290\(00\)00088-5](https://doi.org/10.1016/S0378-4290(00)00088-5)
- Haraguchi, T., Duan, Z., Tamanaha, M., Ito, K., Tominaga, M., 2019. Diversity of Plant Actin–Myosin Systems BT - The Cytoskeleton: Diverse Roles in a Plant's Life. In: Sahi, V.P., Baluška, F. (Eds.), . Springer International Publishing, Cham, pp. 49–61.
- Haraguchi, T., Tominaga, M., Matsumoto, R., Sato, K., Nakano, A., Yamamoto, K., Ito, K., 2014. Molecular characterization and subcellular localization of *Arabidopsis* class VIII myosin, ATM1. *J. Biol. Chem.* 289, 12343–12355. PMID: 24637024.
- Hsiao, Y.-C., Hsu, Y.-F., Chen, Y.-C., Chang, Y.-L., Wang, C.-S., 2016. A WD40 protein, AtGHS40, negatively modulates abscisic acid degrading and signaling genes during seedling growth under high glucose conditions. *J. Plant Res.* 129, 1127–1140. PMID: 27443795.
- Kurth, E.G., Peremyslov, V. V, Turner, H.L., Makarova, K.S., Iranzo, J., Mekhedov, S.L., Koonin, E. V, Dolja, V. V, 2017. Myosin-driven transport network in plants. *Proc. Natl. Acad. Sci.* 114, E1385 LP-E1394. PMID: 28096376.
- Laxmi, A., Paul, L.K., Peters, J.L., Khurana, J.P., 2004. *Arabidopsis* constitutive photomorphogenic mutant, *bls1*, displays altered brassinosteroid response and sugar sensitivity. *Plant Mol. Biol.* 56, 185–201. PMID: 15604737.
- Li, Y., Smith, C., Corke, F., Zheng, L., Merali, Z., Ryden, P., Derbyshire, P., Waldron, K., Bevan, M.W., 2007. Signaling from an altered cell wall to the nucleus mediates sugar-responsive growth and development in *Arabidopsis thaliana*. *Plant Cell* 19, 2500–2515. PMID: 17693536.
- Lilley, J.L.S., Gee, C.W., Sairanen, I., Ljung, K., Nemhauser, J.L., 2012. An endogenous carbon-sensing pathway triggers increased auxin flux and hypocotyl elongation. *Plant Physiol.* 160, 2261–2270. PMID: 23073695.
- Liu, Z., Zhang, Y., Liu, R., Hao, H., Wang, Z., Bi, Y., 2011. Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in *Arabidopsis*. *J. Plant Physiol.* 168, 1771–1779. PMID: 21684034.
- Ma, D., Li, X., Guo, Y., Chu, J., Fang, S., Yan, C., Noel, J.P., Liu, H., 2016. Cryptochrome 1 interacts with PIF4 to regulate high temperature-mediated hypocotyl elongation in response to blue light. *Proc. Natl. Acad. Sci.* 113, 224–229. PMID: 26699514.
- Nakano, T., 2019. Hypocotyl Elongation: A Molecular Mechanism for the First Event in Plant Growth That Influences Its Physiology. *Plant Cell Physiol.* 60, 933–934. PMID: 31050756.
- Oh, S., Warnasooriya, S.N., Montgomery, B.L., 2013. Downstream effectors of light- and phytochrome-dependent regulation of hypocotyl elongation in *Arabidopsis thaliana*. *Plant Mol. Biol.* 81, 627–640. PMID: 23456246.
- Peremyslov, V. V, Klocko, A.L., Fowler, J.E., Dolja, V. V, 2012. *Arabidopsis* Myosin XI-K Localizes to the Motile Endomembrane Vesicles Associated with F-actin. *Front. Plant Sci.* 3, 184. PMID: 22969781.
- Peremyslov, V. V, Prokhnevsky, A.I., Avisar, D., Dolja, V. V, 2008. Two class XI myosins function in organelle trafficking and root hair development in *Arabidopsis*. *Plant Physiol.* 146, 1109–1116. PMID: 18178669.
- Peremyslov, V. V, Prokhnevsky, A.I., Dolja, V. V, 2010. Class XI Myosins Are Required for Development, Cell Expansion, and F-Actin Organization in *Arabidopsis*. *Plant Cell* 22, 1883 LP – 1897. PMID: 20581304.
- Pfeiffer, I., Kutschera, U., 1995. Sucrose metabolism and cell elongation in developing sunflower hypocotyls. *J. Exp. Bot.* 46, 631–638. DOI: <https://doi.org/10.1093/jxb/46.6.631>
- Pollard, T.D., Cooper, J.A., 2009. Actin, a central player in cell shape and movement. *Science* 326, 1208–1212. PMID: 19965462.
- Procko, C., Crenshaw, C.M., Ljung, K., Noel, J.P., Chory, J., 2014. Cotyledon-Generated Auxin Is Required for Shade-Induced Hypocotyl Growth in *Brassica rapa*. *Plant Physiol.* 165, 1285 LP – 1301. PMID: 24891610.
- Prokhnevsky A.I., Peremyslov V.V., Dolja V.V., 2008. Overlapping functions of the four class XI myosins in *Arabidopsis* growth, root hair elongation, and organelle motility. *Proc Natl Acad Sci U S A.* 105(50):19744-19749. PMID: 19060218.
- Reddy, A.S., Day, I.S., 2001. Analysis of the myosins encoded in the recently completed *Arabidopsis thaliana* genome sequence. *Genome Biol.* 2. PMID: 11516337.
- Reed, J.W., Wu, M.-F., Reeves, P.H., Hodgens, C., Yadav, V., Hayes, S., Pierik, R., 2018. Three auxin response factors promote hypocotyl elongation. *Plant Physiol.* 178, 864–875. PMID: 30139794.

- Ryan, J.M., Nebenführ, A., 2018. Update on Myosin Motors: Molecular Mechanisms and Physiological Functions. *Plant Physiol.* 176, 119 LP – 127. PMID: 29162634.
- Salter, M.G., Franklin, K.A., Whitelam, G.C., 2003. Gating of the rapid shade-avoidance response by the circadian clock in plants. *Nature* 426, 680–683. PMID: 14668869.
- Simon, N.M.L., Kusakina, J., Fernández-López, Á., Chembath, A., Belbin, F.E., Dodd, A.N., 2018a. The Energy-Signaling Hub SnRK1 Is Important for Sucrose-Induced Hypocotyl Elongation. *Plant Physiol.* 176, 1299–1310. PMID: 29114081.
- Simon, N.M.L., Sawkins, E., Dodd, A.N., 2018b. Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation. *Plant Signal. Behav.* 13, e1457913. PMID: 29584583.
- Singh, M., Gupta, A., Singh, D., Khurana, J.P., Laxmi, A., 2017. Arabidopsis RSS1 Mediates Cross-Talk Between Glucose and Light Signaling During Hypocotyl Elongation Growth. *Sci. Rep.* 7, 16101. PMID: 29170398.
- Yu, Y., Huang, R., 2017. Integration of ethylene and light signaling affects hypocotyl growth in Arabidopsis. *Front. Plant Sci.* 8, 57. PMID: 28174592.
- Zhang, Y., He, J., 2015. Sugar-induced plant growth is dependent on brassinosteroids. *Plant Signal. Behav.* 10, e1082700–e1082700. PMID: 26340221.
- Zhang, Y., Liu, Z., Wang, J., Chen, Y., Bi, Y., He, J., 2015. Brassinosteroid is required for sugar promotion of hypocotyl elongation in Arabidopsis in darkness. *Planta* 242, 881–893. PMID: 25998528.
- Zhang, Y., Liu, Z., Wang, L., Zheng, S., Xie, J., Bi, Y., 2010. Sucrose-induced hypocotyl elongation of Arabidopsis seedlings in darkness depends on the presence of gibberellins. *J. Plant Physiol.* 167, 1130–1136. PMID: 20430474.
- Zheng, L., Shang, L., Chen, X., Zhang, L., Xia, Y., Smith, C., Bevan, M.W., Li, Y., Jing, H.-C., 2015. TANG1, Encoding a Symplekin\_C Domain-Contained Protein, Influences Sugar Responses in Arabidopsis. *Plant Physiol.* 168, 1000–1012. PMID: 26002908.

**Funding:** This work was supported by an American Association of University Women Research Publication Grant to D.R.K. and Iowa State University start-up funds to D.R.K.

**Author Contributions:** Damilola Olatunji: Conceptualization, Writing - original draft, Writing - review and editing, Formal analysis, Methodology, Investigation, Validation, Visualization. Dior R Kelley: Conceptualization, Writing - original draft, Writing - review and editing, Visualization, Funding acquisition, Resources, Supervision.

**Reviewed By:** Michelle Facette and Anonymous

**History:** Received June 11, 2020 **Revision received** July 3, 2020 **Accepted** July 7, 2020 **Published** July 10, 2020

**Copyright:** © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Olatunji, D; Kelley, DR (2020). A role for Arabidopsis myosins in sugar-induced hypocotyl elongation. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000276>