

Neuronal expression of Ca^{2+} oscillation initiator is linked to rapid neonatal growth in mice

Woojin Kang^{1§}, Kenji Yamatoya², Kenji Miyado¹, Mami Miyado³ and Yoshitaka Miyamoto¹

¹Department of Reproductive Biology, National Research Institute for Child Health and Development, Setagaya, Tokyo 157-8535, Japan

²Institute for Environmental and Gender-Specific Medicine, Juntendo University Graduate School of Medicine, Urayasu, Chiba 279-0021, Japan

³Department of Molecular Endocrinology, National Research Institute for Child Health and Development, Setagaya, Tokyo 157-8535, Japan

[§]To whom correspondence should be addressed: kwjbear@gmail.com

Abstract




Figure 1. Neuronal extra-mitochondrial citrate synthase (eCs) expression and retarded neonatal growth in eCs-deficient (KO) male mice:

(A) Cooperative function of two citrate synthases, CS and eCS. (B) Growth curve of wild-type (WT) and *eCs*-KO mice. Ten mice from each genotype were weighed once a week. Dotted box: *eCs*-KO mice with significantly reduced weight. Arrow: *eCs*-KO mice brains highlighted with *LacZ* staining. Values are expressed as mean \pm standard error of the mean. * $p < 0.0001$. (C) The *eCs* expression in adult mouse brains (9-week-old males). In the *eCs*-null allele, *eCs*-promoter drives the expression of the *LacZ* reporter gene. The *LacZ*-stained brain was viewed from dorsal (upper panel) and ventral (lower panel) surfaces from the same individual male. In the cerebellum, visualization of β -galactosidase activity in KO adult mice showed the *eCs* expression in a single cell layer lined inside the cerebellum, presumably the Purkinje cell layer. Dotted box in the inset: enlarged region. Scale bars: 5 mm.

Description

Repetitive increases in the cytoplasmic calcium concentration (Ca^{2+} oscillation) control a wide variety of biological events (Dupont *et al.* 2011). In fertilization, a sperm-bearing factor, phospholipase C zeta 1 (PLCz1), triggers Ca^{2+} oscillation and resumes cell division in eggs. Citrate synthase (CS) is localized to the mitochondrial matrix, where it catalyzes a reaction between acetyl-coenzyme A (CoA) and oxaloacetate to form citric acid (Surpin and Chory 1997) (Figure 1A). The extra-mitochondrial form of CS (eCS) is encoded by a separate gene in mice and is expressed by alternative splicing from the CS gene in humans (Kang *et al.* 2020). eCS functions as a secondary factor triggering Ca^{2+} oscillation, which is transferred from the sperm to the eggs (Kang *et al.* 2020). More specifically, eCS has been found to trigger an initial Ca^{2+} spike using a PLCz1-independent mechanism. However, the role of eCS-triggered Ca^{2+} oscillation in broad cell functions is unknown.

Presently, we studied the neuronal involvement of eCS and its neuronal expression using eCS-deficient (KO) male mice carrying the *LacZ* reporter gene inserted into the *eCs*-null allele. KO mice were born healthy, but their body size was noticeably small. Therefore, KO mice were weighed daily after birth and compared with wild-type (WT) mice (Figure 1B). From the first to the second week, the body weight was comparable between KO and WT mice. However, KO mice continuously weighed less than WT mice during the third week (6.7 ± 0.5 g vs. 11.2 ± 1.1 g; $p < 0.0001$), fourth week (9.1 ± 0.4 g vs. 14.0 ± 2.5 g; $p < 0.0001$), fifth week (15.6 ± 0.6 g vs. 19.5 ± 0.6 g; $p < 0.0001$), and sixth week (18.3 ± 0.6 g vs. 20.4 ± 0.7 g; $p < 0.0001$). No difference was noted from the seventh week onward. Moreover, eCS expression was detected in a narrow layer of the cerebellar cortex, probably the Purkinje layer (Figure 1C). From this result, we assumed that eCS and eCS-expressed neuronal cells could regulate the rapid increase in body weight during childhood.

Growth retardation is linked to low concentrations of growth hormone (GH) in humans and mice. GH, also known as somatotropin, is a peptide hormone that mainly functions in growth, cell replication, and cell regeneration (Velloso 2008). GH stimulates the production of the insulin-like growth factor-1 (IGF-1) (also called growth-promoting hormone) to regulate overall body growth (Yakar *et al.* 2002). Moreover, GH release conclusively depends on changes in intracellular Ca^{2+} concentration, indicating a critical role of intracellular Ca^{2+} as a mediator of GH function (Cuttler *et al.* 1992). Upon the development of the central nervous system, IGF-1 is abundantly expressed in neurons, especially in Purkinje cells, to promote cell survival in the cerebellum (Torres-Aleman *et al.* 1994; Chrysis *et al.* 2001), suggesting a possible contribution of eCS to neuronal survival.

Otherwise, spontaneous Ca^{2+} oscillation is induced in astrocytes both in vitro and in vivo (Zhou *et al.* 2020). Ca^{2+} oscillation also controls the fate determination of cultured neural stem cells (Glaser *et al.* 2020). From our results, we concluded that eCS expressed in the Purkinje cell layer could trigger neuronal signaling via Ca^{2+} oscillation, subsequently enhancing the rapid growth observed during childhood.

Citrate has been shown to be a regulator of various biological processes, such as insulin secretion (Iacobazzi and Infantino 2014). Citrates act as suppressors of problematic events, such as in the reduction of oxidative stress and inflammation (Iacobazzi and Infantino 2014) and in the protection against traumatic brain injury (Kilbaugh *et al.* 2015). However, while it is known that citrates are specifically synthesized and released from astrocytes (Westergaard *et al.* 2017), presently, knowledge of the role of citrates is insufficient. Our results largely contribute to the understanding of eCS-mediated Ca^{2+} oscillation in the brain.

Methods

[Request a detailed protocol](#)

Animals

Mutant mice were generated from C57BL/6-derived embryonic stem cell clones by injection into blastocysts from C57BL/6 mouse with a genetically deleted *Csl* (*eCs*) (*Csl*^{tm1(KOMP)Vlcg}; ID14519) obtained from the Knockout Mouse Project (KOMP) repository (an NCRR-NIH-supported strain suppository; www.komp.org). Homozygous mice (C57BL/6 genetic background) were generated by subsequent intercrosses of heterozygous animals. For *LacZ* staining, 8-week-old male C57BL/6J mice were purchased from Japan SLC Inc. (Shizuoka, Japan) and their brains were used as control.

All mice were housed under specific, controlled pathogen-free conditions. Food and water were available ad libitum. All animal experiments were approved by The Institutional Animal Care and Use Committee of the National Research Institute for Child Health and Development (Experimental number, A2004-004).

LacZ staining

After fixation with 4% paraformaldehyde (Wako Pure Chemical Industries, Osaka, Japan), mouse brains were washed three times with 0.1 M phosphate-buffered saline (pH 7.4) for 5 min. They were then incubated in β -gal staining solution [1 mg/mL 5-Bromo-4-chloro-3-indolyl β -D-galactopyranoside, 2 mM MgCl_2 , 5 mM potassium hexacyanoferrate (III), and 5 mM potassium hexacyanoferrate (II) trihydrate, 0.01% (w/v) sodium deoxycholate, 0.02 (w/v) NP-40] at 37 °C for overnight. To observe the eCS expression patterns, samples were embedded in Tissue-Tek OCT compound (Sakura,

Finetek, Tokyo, Japan), frozen in liquid nitrogen, and cut into thin sections (10 μm) using a cryostat (CryoStar NX70, Thermo Fisher Scientific, Inc., MA).

Statistical analysis

Comparisons were made using one-way analysis of variance following Scheffe's method, Mann–Whitney *U*-test, or Fisher's exact test. Statistical significance was defined as $p < 0.05$. Results are expressed as the mean \pm standard error of the mean.

Acknowledgments: We greatly appreciate the JAC staff for supporting our experiments through comprehensive animal care services. We also acknowledge Editage (<https://www.editage.jp/>) for English language editing.

References

- Chrysis D, Calikoglu AS, Ye P, D'Ercole AJ. 2001. Insulin-like growth factor-I overexpression attenuates cerebellar apoptosis by altering the expression of Bcl family proteins in a developmentally specific manner. *J Neurosci* 21: 1481-9. DOI: 10.1523/JNEUROSCI.21-05-01481.2001 | PMID: 11222638.
- Cuttler L, Glaum SR, Collins BA, Miller RJ. 1992. Calcium signalling in single growth hormone-releasing factor-responsive pituitary cells. *Endocrinology* 130: 945-53. DOI: 10.1210/endo.130.2.1733736 | PMID: 1733736.
- Dupont G, Combettes L, Bird GS, Putney JW. 2011. Calcium oscillations. *Cold Spring Harb Perspect Biol* 3: . DOI: 10.1101/cshperspect.a004226 | PMID: 21421924.
- Glaser T, Shimojo H, Ribeiro DE, Martins PPL, Beco RP, Kosinski M, Sampaio VFA, Corrêa-Velloso J, Oliveira-Giacomelli Á, Lameu C, de Jesus Santos AP, de Souza HDN, Teng YD, Kageyama R, Ulrich H. 2020. ATP and spontaneous calcium oscillations control neural stem cell fate determination in Huntington's disease: a novel approach for cell clock research. *Mol Psychiatry* : . DOI: 10.1038/s41380-020-0717-5 | PMID: 32350390.
- Iacobazzi V, Infantino V. 2014. Citrate--new functions for an old metabolite. *Biol Chem* 395: 387-99. DOI: 10.1515/hsz-2013-0271 | PMID: 24445237.
- Kang W, Harada Y, Yamatoya K, Kawano N, Kanai S, Miyamoto Y, Nakamura A, Miyado M, Hayashi Y, Kuroki Y, Saito H, Iwao Y, Umezawa A, Miyado K. 2020. Extra-mitochondrial citrate synthase initiates calcium oscillation and suppresses age-dependent sperm dysfunction. *Lab Invest* 100: 583-595. DOI: 10.1038/s41374-019-0353-3 | PMID: 31857692.
- Kilbaugh TJ, Karlsson M, Byro M, Bebee A, Ralston J, Sullivan S, Duhaime AC, Hansson MJ, Elmer E, Margulies SS. 2015. Mitochondrial bioenergetic alterations after focal traumatic brain injury in the immature brain. *Exp Neurol* 271: 136-44. DOI: 10.1016/j.expneurol.2015.05.009 | PMID: 26028309.
- Surpin M, Chory J. 1997. The co-ordination of nuclear and organellar genome expression in eukaryotic cells. *Essays Biochem* 32: 113-25. PMID: 9493015.
- Torres-Aleman I, Pons S, Arévalo MA. 1994. The insulin-like growth factor I system in the rat cerebellum: developmental regulation and role in neuronal survival and differentiation. *J Neurosci Res* 39: 117-26. DOI: 10.1002/jnr.490390202 | PMID: 7530775.
- Velloso CP. 2008. Regulation of muscle mass by growth hormone and IGF-I. *Br J Pharmacol* 154: 557-68. DOI: 10.1038/bjp.2008.153 | PMID: 18500379.
- Westergaard N, Waagepetersen HS, Belhage B, Schousboe A. 2017. Citrate, a Ubiquitous Key Metabolite with Regulatory Function in the CNS. *Neurochem Res* 42: 1583-1588. DOI: 10.1007/s11064-016-2159-7 | PMID: 28058526.
- Yakar S, Rosen CJ, Beamer WG, Ackert-Bicknell CL, Wu Y, Liu JL, Ooi GT, Setser J, Frystyk J, Boisclair YR, LeRoith D. 2002. Circulating levels of IGF-1 directly regulate bone growth and density. *J Clin Invest* 110: 771-81. DOI: 10.1172/JCI15463 | PMID: 12235108.
- Zhou A, Liu X, Zhang S, Huo B. 2020. Effects of store-operated and receptor-operated calcium channels on synchronization of calcium oscillations in astrocytes. *Biosystems* 198: 104233. DOI: 10.1016/j.biosystems.2020.104233 | PMID: 32858094.

Funding: This work was supported in part by JSPS KAKENHI, Grant Numbers JP19H01067 and JP19K09793.

Author Contributions: Woojin Kang: Supervision, Writing - review and editing. Kenji Yamatoya: Investigation. Kenji Miyado: Writing - original draft, Writing - review and editing. Mami Miyado: Investigation. Yoshitaka Miyamoto: Investigation.

Reviewed By: Anonymous

History: Received October 8, 2020 **Revision received** November 10, 2020 **Accepted** November 10, 2020 **Published** November 12, 2020

Copyright: © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Kang, W; Yamatoya, K; Miyado, K; Miyado, M; Miyamoto, Y (2020). Neuronal expression of Ca²⁺ oscillation initiator is linked to rapid neonatal growth in mice. microPublication Biology. <https://doi.org/10.17912/micropub.biology.000325>