

# The conserved MASRPF motif in the Attractin homolog, Distracted, is required for association with *Drosophila* E3-ligase Mgrn1

Vindhya Nawaratne<sup>1</sup>, Sirisha Kudumala<sup>2</sup>, Priyanka Prakash Kakad<sup>3</sup> and Tanja A Godenschwege<sup>1§</sup>

<sup>1</sup>Biological Science Department, Florida Atlantic University, Jupiter, FL 33458


<sup>2</sup>Prodigy Biotech, Jupiter, FL 33458

<sup>3</sup>ScienCell Research Laboratories, Inc., 1610 Faraday Avenue, Carlsbad, CA 92008

<sup>§</sup>To whom correspondence should be addressed: godensch@fau.edu

## Abstract

In rodents, all three paralogs of the Attractin (Atrn) transmembrane protein family exhibit strong phenotypic overlap and are implicated in the regulation of the same G-protein coupled receptors (GPCR) as E3-ligase Mahogunin ring finger 1 (Mgrn1). Recently it was shown that the highly conserved intracellular MASRPF motif in mammal Multiple epidermal growth factor-like domain 8 protein is required for binding of Mgrn1 to mediate ubiquitination of GPCR Smoothed in vitro. Here, we show that the MASRPF motif of *Drosophila* Distracted, the ortholog of ATRN and Attractin-like 1, is required for association with *Drosophila* Mgrn1 (dMgrn1) in vivo.



**Figure 1. In vivo interaction of Distracted and dMgrn1** : **A**. Comparison of protein motifs of *Drosophila* and human Attractin (ATRNL1) family orthologs. In mice, the MASRPF motif of Multiple EGF like domains 8 (Megf8) was shown to be essential for Mgrn1 binding (Kong *et al.*, 2020). Here we show that *Drosophila* CG9941 (dMgrn1) associates with Dsd in a MASRPF motif-dependent manner. **B**. Sequence alignment of the first 40 membrane-proximal amino acids in the intracellular domain of *Drosophila* and human ATRNL1 orthologs. In *Drosophila*, the MASRPF motif (blue) is fully conserved in distracted (*dsd*), but only partially in the *Megf8* gene. Bar indicates amino acids deleted in the myc-tagged *Dsd*ΔMASR construct. **C**. Co-immunoprecipitation of Dsd and dMgrn1. Top blot was probed with anti-FLAG antibodies for dMgrn1, and subsequently probed with anti-myc antibodies for Dsd-myc IP (bottom blot). First three lanes show 5% of the input of protein homogenates obtained from *Drosophila* heads that were panneuronally (*nSyb-Gal4* driver) expressing FLAG-tagged *UAS-dMgrn1* (lane 1), co-expressing *UAS-dMgrn1-FLAG* with *UAS-Dsd*ΔMASR-myc (lane 2) or *UAS-Dsd-myc* (lane 3). Lanes 3-6 show immunoprecipitation (IP) for myc-tagged proteins of the supernatant containing only dMgrn1-FLAG (lane 4, negative control), dMgrn1-FLAG and DsdΔMASR-myc (lane 5), or dMgrn1-FLAG and wildtype (WT) Dsd-myc (lane 6). dMgrn1-FLAG is detected at approximately 110kDa and Dsd-myc at 160kDa. Asterisk indicates an unspecific protein detected by the antibodies.

## Description

In vertebrates, the Attractin (Atrn) transmembrane protein family is comprised of ATRNL1, Attractin-like 1 (ATRNL1) and Multiple epidermal growth factor-like domain 8 (MEGF8) proteins, which contain an extensive extracellular domain with numerous protein interaction motifs and a highly conserved intracellular MASRPF motif (Fig. 1A and B) (Walker *et al.*,

2007). Common to all three paralogs is that they have a strong phenotypic overlap with E3-ligase Mahogunin ring finger 1 (Mgmn1), due to regulation of the same G-protein coupled receptors (GPCRs) as Mgmn1. For example, rodent *Atrn* and *Mgmn1* negatively regulate signaling melanocortin receptors MCR1 and MCR4 by promoting their endolysosomal trafficking. Thus, loss of function mutations in *Atrn* or *Mgmn1* suppress the pigmentation and obesity phenotypes of MCR1/4 antagonist gain of function mutants (Barsh *et al.*, 2002; He *et al.*, 2003; Nagle *et al.*, 1999). Loss of function of *Megf8* and *Mgmn1* mutants exhibit common developmental phenotypes (Aune *et al.*, 2008; Cota *et al.*, 2006; Zhang *et al.*, 2009) and both proteins were identified as negative regulators of the GPCR Smoothed in mice (Pusapati *et al.*, 2018). Most recently it was shown in vitro that *Mgmn1* and *Megf8* co-immunoprecipitate in a MASRPF motif-dependent manner, and this association is required for *Mgmn1* to mediate membrane-tethered ubiquitination to modulate the signaling strength of Hedgehog morphogens in mice (Kong *et al.*, 2020).

Distracted (*dsd*) is the sole *Drosophila* ortholog of human ATRN and ATRNL1 (Fig. 1A), while *Drosophila* *Megf8*, like rodent *Megf8*, is required for early development (Lloyd *et al.*, 2018). In contrast to *Drosophila* *Megf8*, the MASRPF motif is fully conserved in *Dsd* (Fig. 1B). The DRSC Integrative Ortholog Prediction Tool uses 15 different algorithms to identify orthologous genes ([https://www.flyrnai.org/cgi-bin/DRSC\\_orthologs.pl](https://www.flyrnai.org/cgi-bin/DRSC_orthologs.pl), Version 8.0, August 2019). For *CG9941* (hereafter referred to as *dMgmn1*) more algorithms predicted that it is orthologous to mammal *MGRN1* (human 11/15, rat 7/13, mouse 10/15) than for *RNF157* (Ring Finger Protein 157; human 10/14, rat 4/15, mouse 10/15). While the zinc finger domain is highly conserved (83% amino acid identity) in *dMgmn1* when compared to either mammal *MGRN1* or *RNF157* proteins, the entire protein is less well preserved (amino acid identity 35-37%/similarity 48-51%) in part due to the *Drosophila* protein being larger by approximately 210 amino acids when compared to mammalian *Mgmn1*.

Here, we generated myc-tagged *Dsd* containing (*UAS-Dsd-myc*) or lacking the MASRPF motif (*UAS-DsdΔMASR-myc*, Fig. 1B), and FLAG-tagged *dMgmn1* (*UAS-dMgmn1-FLAG*) transgenic lines to determine whether *Dsd* and *dMgmn1* are in a complex together in vivo, and if the interaction is dependent on the MASRPF motif. We panneuronally (*nSyb-Gal4*) expressed *dMgmn1-FLAG* alone or together with *Dsd-myc* or *DsdΔMASR-myc*, and used *Drosophila* heads to perform immunoprecipitation (IP) with myc-trap agarose beads. In western blots of fly head supernatant prior to IP, *dMgmn1-FLAG* is detected at 110 kDa and *Dsd-myc* at 160kDa (Fig. 1, lane 1-3) confirming protein expression. Multiple independent IP experiments were performed using two distinct transgenic lines for expression of *Dsd-myc* and *DsdΔMASR-myc*, while using the same transgenic line for expression of *dMgmn1-FLAG*. The results demonstrated that *Dsd* and *dMgmn1* co-IP, suggesting they interact in vivo (n=4), and that this interaction was indeed dependent on the MASRPF motif (n=3, Fig. 1C). After co-IP, *dMgmn1-FLAG* protein was detected abundantly when it was co-expressed with *Dsd-myc* (Fig. 1C, lane 6) but absent or at similarly low levels as the negative controls (*UAS-dMgmn1-FLAG,nSyb-Gal4*, Fig. 1C, lane 4) when co-expressed with *DsdΔMASR-myc* (Fig. 1C, lane 5). These results imply that *Drosophila* is a valid model to study ATRN's proposed role as a substrate adaptor for MGRN1 (Kong *et al.*, 2020). Loss of function of *Atrn* and *Mgmn1* in rodents is associated with mitochondrial dysfunction, elevated oxidative stress, and adult-onset spongiform neurodegeneration (Bronson *et al.*, 2001; Cota *et al.*, 2008; He *et al.*, 2003; Kuramoto *et al.*, 2001; Li *et al.*, 2014; Muto and Sato, 2003; Paz *et al.*, 2007). In humans, ATRN is a candidate gene for sporadic amyotrophic lateral sclerosis (Morahan *et al.*, 2009), ATRN plasma levels are altered in early-onset Alzheimer's disease (AD) patients prior to symptoms (Muenchhoff *et al.*, 2016), and its gene expression is reduced in Parkinson disease (PD) patients (Glaab and Schneider, 2015). Since the ATRN/MGRN1 receptor substrates involved in nervous system survival have not been identified, future studies in *Drosophila* may provide insight into the relevant signaling pathways.

## Methods

[Request a detailed protocol](#)

### Cloning and mutagenesis of *Dsd* and *dMgmn1* transgenes

All primers were generated by Integrated DNA Technologies (IDT, Iowa, USA). The *Dsd* open reading frame (ORF) without the stop codons was amplified with forward (CAC CAT GTC CCT GTT GCC ACC G) and reverse (TGT ACA ACT GTC TGG GTG CTG GCT CTG) primers from the LD14047 cDNA clone (Stock Number: 1301485, *Drosophila* Genomics Resource Center, Indiana, USA) and cloned into the pENTR/D-TOPO (Invitrogen, Massachusetts, USA). The *dMgmn1* open reading frame without the stop codons was amplified with forward (ATG GGC AAC TTG TGA GCA G) and reverse (AAC ATT GAC GGC ATT CTG) primers from the LP12254 cDNA clone (Stock Number: 1344980, *Drosophila* Genomics Resource Center, Indiana, USA) and cloned into the pENTR/D-TOPO. In addition, the highly conserved motif MASRPFSQVLV (amino acids 1208-1218) was deleted in pENTR/D-TOPO-DSD entry clone with QuikChange Lightning Site-Directed Mutagenesis Kit (Agilent, California, USA) using forward (CGT CGA AAT GGA ACA GGA TAT CGA GAA CCG CG) and reverse (CGC GGT TCT CGA TAT CCT GTT CCA TTT CGA CG) primers. The vectors pENTR/D-TOPO containing *Mgmn1*, *Dsd* without or with the MASRPFSQVLV deletion (referred to as *DsdΔMASR*) were verified using restriction enzyme digests and Sanger sequencing (Genewiz, New Jersey, USA). Both *Dsd* entry vectors were recombined with the pTWM destination vector (Stock Number: 1107, *Drosophila* Genomics Resource Center, Indiana, USA) and the *dMgmn1* entry vector with the pTWF destination vector (Stock Number: 1116, *Drosophila* Genomics Resource Center, Indiana, USA) using LR clonase. The recombination for the presence of six in-

frame myc-codons for the *Dsd* constructs and the three in-frame FLAG codons for the *Mgrn1* construct at the C-termini was verified using restriction enzyme digests and Sanger sequencing (Genewiz, New Jersey, USA). The multiple UAS-transgenic lines with P-element insertions at random sites on the second and third chromosome for all three constructs were established in the *w<sup>1118</sup>* background by BestGene Inc (California, USA).

#### *Fly strains and maintenance*

The *nSyb-Gal4* line (Bloomington Stock Center, Indiana, USA, RRID: BDSC\_51635) was recombined with the *UAS-dMgrn1-FLAG* line to establish a stable stock expressing dMgrn1-FLAG panneuronally. The *UAS-dMgrn1-FLAG,nSyb-Gal4* stock was crossed to two distinct *UAS-Dsd-myc* or the *UAS-DsdΔMASR-myc* lines and the offspring were collected for the co-IP experiments. The *UAS-dMgrn1-FLAG,nSyb-Gal4* stock was used as a negative control. All flies were reared on standard fly media at 25°C.

#### *Co-immunoprecipitation*

Collected adult flies (~3ml) were snap-frozen in liquid nitrogen in 15ml conical tubes, heads were severed by vigorous shaking, and collected by passing through a sieve. Samples were kept at 4°C for the rest of the experiment. 100mg of heads were homogenized in 450µl of lysis buffer (10 mM Tris base, 150 mM NaCl, 0.5 mM EDTA, 0.5% NP-40, 10µM MG132, 1 mM PMSF, 20mM NEM, 1x Complete ULTRA EDTA-free Protease Inhibitor cocktail, 1xPhosStop) using a handheld homogenizer. Homogenates were centrifuged at 500xg for 1min to remove cuticle particulates, and the supernatant was collected and centrifuged again at 16000xg for 30min. 5% of the supernatant was saved as inputs while the rest of the supernatant was precleared with 60µl of Protein A agarose beads (Pierce™ Protein A Agarose, ThermoFisher Scientific, Massachusetts, USA) for 10min. Protein A agarose beads were removed by centrifugation at 2500xg for 2 min and supernatant immunoprecipitated with 25µl of Myc-Trap® (ChromoTek, Bavaria, Germany) agarose beads for 1hr. Beads were collected by centrifugation at 2500xg for 2 min, washed 3 times with 500µl dilution buffer (10mM Tris base, 150mM NaCl), resuspended in 30µl 2x Laemmli buffer, boiled for 10 min at 95°C and pelleted at 2500xg for 2 min. Proteins in the supernatant were separated on an 8% SDS-PAGE gel. Western blots of the gels were first probed with mouse anti-FLAG® M2 (Sigma-Aldrich, Missouri, USA, RRID:AB\_259529, dilution 1:1000) and with mouse anti-myc 9E11 antibody (BioLegend, California, USA, RRID: AB\_2565039, dilution 1:1000) consecutively. Goat anti-mouse IRDye® 680RD (LI-COR Biosciences, Nebraska, USA, RRID:AB\_10956588, dilution 1:15000) was used to detect the primary antibodies. Images were acquired and analyzed using Image Studio from LI-COR Biosciences.

**Acknowledgments:** Destination vectors and cDNA clone were obtained from Drosophila Genomics Resource Center (NIH Grant 2P40OD010949) and the *nSyb-Gal4* line from the Bloomington Drosophila Stock Center (Indiana, United States, NIH P40OD018537).

## References

- Alberts MJ. 2001. Suppression of recurrent transient ischemic attacks by a statin agent. *Neurology* 56: 531-2. PMID: 11222801.
- Aune CN, Chatterjee B, Zhao XQ, Francis R, Bracero L, Yu Q, Rosenthal J, Leatherbury L, Lo CW. 2008. Mouse model of heterotaxy with single ventricle spectrum of cardiac anomalies. *Pediatr Res* 63: 9-14. PMID: 18043505.
- Barsh GS, He L, Gunn TM. 2002. Genetic and biochemical studies of the Agouti-attractin system. *J Recept Signal Transduct Res* 22: 63-77. PMID: 12503608.
- Cota CD, Bagher P, Pelc P, Smith CO, Bodner CR, Gunn TM. 2006. Mice with mutations in Mahogunin ring finger-1 (*Mgrn1*) exhibit abnormal patterning of the left-right axis. *Dev Dyn* 235: 3438-47. PMID: 17075880.
- Cota CD, Liu RR, Sumberac TM, Jung S, Vencato D, Millet YH, Gunn TM. 2008. Genetic and phenotypic studies of the dark-like mutant mouse. *Genesis* 46: 562-73. PMID: 18821597.
- Glaab E, Schneider R. 2015. Comparative pathway and network analysis of brain transcriptome changes during adult aging and in Parkinson's disease. *Neurobiol Dis* 74: 1-13. PMID: 25447234.
- He L, Eldridge AG, Jackson PK, Gunn TM, Barsh GS. 2003. Accessory proteins for melanocortin signaling: attractin and mahogunin. *Ann N Y Acad Sci* 994: 288-98. PMID: 12851328.
- He L, Lu XY, Jolly AF, Eldridge AG, Watson SJ, Jackson PK, Barsh GS, Gunn TM. 2003. Spongiform degeneration in mahoganoid mutant mice. *Science* 299: 710-2. PMID: 12560552.
- Kong JH, Young CB, Pusapati GV, Patel CB, Ho S, Krishnan A, Lin JI, Devine W, Moreau de Bellaing A, Athni TS, Aravind L, Gunn TM, Lo CW, Rohatgi R. 2020. A Membrane-Tethered Ubiquitination Pathway Regulates Hedgehog Signaling and Heart Development. *Dev Cell* 55: 432-449.e12. PMID: 32966817.
- Kuramoto T, Kitada K, Inui T, Sasaki Y, Ito K, Hase T, Kawaguchi S, Ogawa Y, Nakao K, Barsh GS, Nagao M, Ushijima T, Serikawa T. 2001. Attractin/mahogany/zitter plays a critical role in myelination of the central nervous system. *Proc Natl*

Acad Sci U S A 98: 559-64. PMID: 11209055.

Li J, Yang J, Cheng D, Shen SL, Xiong CL. 2014. New clues to identify proteins correlated with Attractin. *Andrologia* 46: 796-804. PMID: 23998293.

Lloyd DL, Toegel M, Fulga TA, Wilkie AOM. 2018. The *Drosophila* homologue of MEGF8 is essential for early development. *Sci Rep* 8: 8790. PMID: 29884872.

Morahan JM, Yu B, Trent RJ, Pamphlett R. 2009. A genome-wide analysis of brain DNA methylation identifies new candidate genes for sporadic amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 10: 418-29. PMID: 19922134.

Muenchhoff J, Poljak A, Thalamuthu A, Gupta VB, Chatterjee P, Raftery M, Masters CL, Morris JC, Bateman RJ, Fagan AM, Martins RN, Sachdev PS. 2016. Changes in the plasma proteome at asymptomatic and symptomatic stages of autosomal dominant Alzheimer's disease. *Sci Rep* 6: 29078. PMID: 27381087.

Muto Y, Sato K. 2003. Pivotal role of attractin in cell survival under oxidative stress in the zitter rat brain with genetic spongiform encephalopathy. *Brain Res Mol Brain Res* 111: 111-22. PMID: 12654511.

Nagle DL, McGrail SH, Vitale J, Woolf EA, Dussault BJ Jr, DiRocco L, Holmgren L, Montagno J, Bork P, Huszar D, Fairchild-Huntress V, Ge P, Keilty J, Ebeling C, Baldini L, Gilchrist J, Burn P, Carlson GA, Moore KJ. 1999. The mahogany protein is a receptor involved in suppression of obesity. *Nature* 398: 148-52. PMID: 10086355.

Paz J, Yao H, Lim HS, Lu XY, Zhang W. 2007. The neuroprotective role of attractin in neurodegeneration. *Neurobiol Aging* 28: 1446-56. PMID: 16860906.

Pusapati GV, Kong JH, Patel BB, Krishnan A, Sagner A, Kinnebrew M, Briscoe J, Aravind L, Rohatgi R. 2018. CRISPR Screens Uncover Genes that Regulate Target Cell Sensitivity to the Morphogen Sonic Hedgehog. *Dev Cell* 44: 271. PMID: 29401421.

Walker WP, Aradhya S, Hu CL, Shen S, Zhang W, Azarani A, Lu X, Barsh GS, Gunn TM. 2007. Genetic analysis of attractin homologs. *Genesis* 45: 744-56. PMID: 18064672.

Zhang Z, Alpert D, Francis R, Chatterjee B, Yu Q, Tansey T, Sabol SL, Cui C, Bai Y, Koriabine M, Yoshinaga Y, Cheng JF, Chen F, Martin J, Schackwitz W, Gunn TM, Kramer KL, De Jong PJ, Pennacchio LA, Lo CW. 2009. Massively parallel sequencing identifies the gene *Megf8* with ENU-induced mutation causing heterotaxy. *Proc Natl Acad Sci U S A* 106: 3219-24. PMID: 19218456.

**Funding:** The project was supported by the Jupiter Life Sciences Initiative and the Brain Institute pilot grant at Florida Atlantic University and the National Institute for Neurological Disease and Stroke (R15NS090043)

**Author Contributions:** Vindhya Nawaratne: Methodology, Investigation, Validation, Writing - original draft, Writing - review and editing, Formal analysis. Sirisha Kudumala: Investigation, Writing - review and editing, Validation. Priyanka Prakash Kakad: Writing - review and editing, Supervision, Investigation, Validation. Tanja A Godenschwege: Conceptualization, Funding acquisition, Supervision, Visualization, Writing - original draft, Writing - review and editing.

**Reviewed By:** Anonymous

**History:** Received March 27, 2021 **Revision received** June 14, 2021 **Accepted** June 15, 2021 **Published** July 2, 2021

**Copyright:** © 2021 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Nawaratne, V; Kudumala, S; Kakad, PP; Godenschwege, TA (2021). The conserved MASRPF motif in the Attractin homolog, Distracted, is required for association with *Drosophila* E3-ligase Mgrn1. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000416>