

Ecdysone and 20-hydroxyecdysone are not required to activate glycolytic gene expression in *Drosophila melanogaster* embryos

Jason M. Tennessen^{1§}

¹Indiana University

[§]To whom correspondence should be addressed: jtenness@iu.edu

Abstract

Many of the *Drosophila* enzymes involved in carbohydrate metabolism are coordinately up-regulated approximately midway through embryogenesis. Previous studies have demonstrated that this metabolic transition is controlled by the *Drosophila* Estrogen-Related Receptor (dERR), which is stabilized and activated immediately prior to onset of glycolytic gene expression. The mechanisms that promote dERR activity, however, are poorly understood and other transcriptional regulators could control this metabolic transition, independent of dERR. In this regard, the steroid hormone 20-hydroxyecdysone (20E) represents an intriguing candidate for regulating glycolytic gene expression in embryos – not only does the embryonic 20E pulse immediately precede transcriptional up-regulation of glycolytic metabolism, but 20E is also known to promote *Lactate dehydrogenase* gene expression. Here I test the hypothesis that embryonic 20E signaling is required to activate glycolytic gene expression. Using developmental northern blots, I demonstrate that the transcriptional up-regulation of glycolytic genes during embryogenesis still occurs in *shadow* mutants, which are unable to synthesize either ecdysone or 20E. My finding indicates that ecdysone and 20E signaling are not required for this mid-embryonic metabolic transition.




Figure 1. Timecourse northern blots examining glycolytic gene expression in control and *shadow* mutant embryos: Total RNA from stage *w¹¹¹⁸* control embryos and *w¹¹¹⁸; shadow^{10D104/5F45}* mutant embryos were analyzed by northern blot hybridization to detect transcripts encoding DHR3, Pgi, Pfk, Pglym78, and Ldh. The Ldh short exposure represents a 6-hour

long film exposure. The *Ldh* long exposure is the same blot exposed for 24 hours. Hybridization to detect *rp49* mRNA is included as a loading control.

Description

During the course of the *Drosophila* development, metabolism readily adapts to meet the energetic and biosynthetic demands of each life stage (Gillette *et al.*, 2021). This relationship between growth and metabolism is particularly apparent when examining changes in glycolytic gene expression. Approximately midway through embryonic development, transcripts representing nearly every enzyme involved in glycolysis are coordinately up-regulated (Abu-Shumays and Fristrom, 1997; Currie and Sullivan, 1994a, b; Madhavan *et al.*, 1972; Roselli-Reh fuss *et al.*, 1992; Shaw-Lee *et al.*, 1992; Shaw-Lee *et al.*, 1991; Sun *et al.*, 1988; Tennessen *et al.*, 2011; Tennessen *et al.*, 2014; Tixier *et al.*, 2013; Wright and Shaw, 1970). The resulting glycolytic program is maintained throughout the larval growth period and subsequently down-regulated prior to the onset of metamorphosis (White *et al.*, 1999). These metabolic transitions are highly predictable and provide an opportunity to understand how metabolism adapts to meet the energetic demands of insect development.

Activation of glycolytic metabolism in *Drosophila* embryos depends on the *Drosophila* Estrogen-Related Receptor (dERR; FBgn0035849), which represents the sole fly ortholog of the orphan class of ERR nuclear receptors (Ostberg *et al.*, 2003). In embryos cultured at 25°C, dERR becomes transcriptionally active ~10-12 hours after oviposition, resulting in the coordinate up-regulation of genes involved in carbohydrate metabolism (Tennessen *et al.*, 2011). The mechanisms that temporally regulate dERR activity during this metabolic switch, however, remain unknown and additional factors could be controlling embryonic glycolytic metabolism independent of dERR. In this regard, both dERR activation and onset of glycolytic gene expression correlate with the embryonic pulse of the steroid hormone 20-hydroxyecdysone (20E), which triggers dorsal closure, cuticle deposition, head involution, and a variety of other developmental events (Chavez *et al.*, 2000; Kozlova and Thummel, 2003; Maróy *et al.*, 1988; Warren *et al.*, 2002). Moreover, not only is 20E known to induce *Lactate Dehydrogenase* (*Ldh*; FBgn0001258) expression in imaginal discs (Abu-Shumays and Fristrom, 1997), but the *Ecdysone Receptor* (*EcR*; FBgn0000546) and dERR are also reported to cooperatively regulate expression of glycolytic genes in insect cell culture (Kovalenko *et al.*, 2019), thus raising the possibility that 20E-signaling is necessary for the up-regulation of embryonic carbohydrate metabolism.

To test the hypothesis that 20E transcriptionally regulates glycolysis in *Drosophila* embryos, I used developmental northern blots to analyze expression of key glycolytic genes in *shadow* (*sad*; FBgn0003312) mutants, which are unable to synthesize either ecdysone or 20E (Warren *et al.*, 2002). As a positive control, I first analyzed expression of *DHR3* (FBgn0000448), which is up-regulated in response to embryonic 20E production (Ruaud *et al.*, 2010). Unlike *w¹¹¹⁸* control samples that expressed *DHR3* during mid-embryogenesis, *DHR3* transcripts were undetectable in *sad* mutant samples – a result consistent with lack of 20E production in these embryos. In contrast, *sad* mutants still exhibit the coordinate up-regulation of *Phosphoglucosomerase* (*Pgi*; FBgn0003074), *Phosphofructokinase* (*Pfk*; FBgn0003071), and *Phosphoglyceromutase 78* (*Pgym78*; FBgn0014869) transcripts during mid-embryogenesis, indicating that 20E is not necessary to activate glycolytic gene expression during mid-embryogenesis. I would note, however, that *Ldh* transcripts levels were noticeably decreased at the 10-12 hour AEL timepoint in *sad* mutant samples when compared with controls, suggesting that 20E partially regulates *Ldh* expression at this early timepoint. Such a possibility is consistent with previous observations that 20E activates *Ldh* expression in imaginal discs and that embryonic *Ldh* mRNA transcript levels are elevated at an earlier timepoint when compared with other glycolytic genes (Abu-Shumays and Fristrom, 1997; Tennessen *et al.*, 2011).

Overall, my findings suggest that that 20E signaling is not required for the coordinate up-regulation of glycolytic gene expression observed midway through embryonic development. All glycolytic genes examined in this study exhibited normal up-regulation in *sad* mutants, with the only exception being *Ldh*, which exhibited slightly delayed onset of expression when compared with the control strain. My finding, however, does not exclude the possibility that ecdysone or 20E controls glycolytic metabolism within individual cells or tissues that would be overlooked using northern blot analysis. Finally, my study further highlights the role of dERR as a central regulator of carbohydrate metabolism during mid-embryogenesis and again raises the question as to what developmental signals induce dERR activity during this metabolic transition.

Methods

[Request a detailed protocol](#)

Drosophila Genetics and Embryo Collection: All strains were maintained on Bloomington *Drosophila* Stock Center (BDSC) media. Strains containing the *sad* alleles *sad^{10D104}* and *sad^{5F45}* (a kind gift from Dr. Michael O'Connor) were individually crossed to BDSC stock #6663 (*w¹¹¹⁸*; *Dr^{Mio}/TM3*, *P{w[+mC]=GAL4-twi.G}2.3*, *P{UAS-2xEGFP}AH2.3*, *Sb¹ Ser¹*). F1 male

offspring were again crossed into BDSC strain #6663 and progeny lacking the *Dr^{Mio}* chromosome were used to establish balanced stocks.

Egg collection and synchronization were conducted as previously described (Li and Tennessen, 2017). For both *w¹¹¹⁸* controls and *sad* mutants, 50 females and 25 males were placed in an egg-laying bottle with a molasses agar plate containing a smear of yeast paste taped in the lid. In the case of *sad* mutant embryo collections, *sad^{10D104/TM3}*, *P{w[+mC]=GAL4-twi.G}2.3*, *P{UAS-2xEGFP}AH2.3*, *Sb¹ Ser¹* virgin females were crossed with *sad^{5F45/TM3}*, *P{w[+mC]=GAL4-twi.G}2.3*, *P{UAS-2xEGFP}AH2.3*, *Sb¹ Ser¹* males. *sad^{10D104/5F45}* mutant embryos were identified by the absence of GFP expression using a Zeiss SterEO Discovery V8 microscope.

Northern Blots: RNA preparation and northern blot analysis was conducted using a previously described method (Karim and Thummel, 1992). Briefly, staged embryos were dechorionated and RNA extracted using Trizol Reagent (Life Technologies) following the manufacturers protocol. 3 μg total RNA from each sample was individually added to a 1.5 ml microcentrifuge tube containing a premixed solution of 3 μl 10X Formaldehyde Gel Buffer (0.2 M MOPS pH 7.0, 50 mM Sodium Acetate, 10 mM EDTA), 3.5 μl of formaldehyde (37% w/v), and 10 μl of formamide. Sample volume was adjusted to 25 μl by adding the appropriate volume of nuclease free H₂O, incubated at 65°C for 5 minutes, and centrifuged briefly at 10,000 x g. 3 μl of loading dye (80% glycerol, 1 mM EDTA, 0.4% bromophenol blue, 0.4% xylene cyanol) was added to the sample and mixed by pipetting. The entire sample was then loaded into a formaldehyde-agarose gel (1 g agarose, 75 mL H₂O, 10 mL of 10X formaldehyde gel buffer, 15 mL of 37% formaldehyde [m/v]; immersed within 1X formaldehyde gel buffer). RNA samples were separated at 70 volts for 150 minutes.

RNA was transferred overnight from the gel to GeneScreen Hybridization Transfer Membrane (Perkin Elmer) using standard blotting techniques and crosslinked to the membrane using the “Autocrosslink” setting on a Stratagen UV Stratalinker 1800. Prior to hybridization, membranes were pretreated for one hour at 42°C with 10 ml of hybridization buffer (5 ml of formamide, 2 ml of 10X PIPES buffer [0.1 M PIPES, pH6.5, 4 M NaCl], 1 ml of 10% SDS, 2 ml H₂O, 100 μl sheared herring sperm DNA).

For Pfk, Pgi, Pglym78, and Ldh, radioactive probes were generated from PCR amplified cDNA fragments (see reagents table for oligos used to synthesize cDNA fragments). For DHR3 and rp49, probes were generated as previously described (Sullivan and Thummel, 2003). Labeling reactions were conducted with ³²P-labeled dCTP ([α-³²P]- 3000Ci/mmol 10mCi/ml EasyTide, 250 μC; Perkin Elmer; BLU513H250UC) using a Prime-It II Random Primer Labeling Kit (Agilent Catalog #300385) following the manufacturer’s instructions. Individual reactions were cleaned using a MicroSpin G-50 Sephadex column (Amersham Catalog #27533001). Radioactive probes were added to 100 μl of sheared herring sperm DNA in a 2 ml screwcap tube, boiled for 5 minutes, and added to the hybridization tube containing the membrane and buffer.

Labeled probes were allowed to hybridize with the membrane overnight at 42°C, at which time the hybridization buffer/probe mixture was poured out of the hybridization tube and the membrane washed twice for 10 minutes at 42°C with 10 ml of 2X SSC + 0.1% SDS, once for 10 min at 55°C with 10 ml of 1X SSC + 0.1% SDS, and once for 10 min at 55°C with 10 ml of 0.1X SSC + 0.1% SDS. After the final wash, the membrane was removed from the hybridization tube, briefly immersed in 2X SSC, wrapped in plastic wrap, sandwiched between two intensifying screens with a piece of film, and placed at -80°C for 24 hours. Exposed film was developed using a Kodak X-OMAT film processor.

Reagents

Oligo sequences used to generate northern blot probes

Gene	Oligo sequences
Pfk	5'- ATGCATTCAATAAAATTTTCGAGTATTTACC-3' 5'- TTAGGCGACGGCGTCAGTGTCAC-3'
Pgi	5'- ATGGCCGGCCCACTTCCTCC -3' 5'- TTACTTCCAATTGGCTTTGATG-3'
Pglym78	5'- CCACTACGGTGGACTCACTG -3' 5'- ATGGCCTTCTTCACGGTCTC -3'

Ldh	5'- ATGCCCGCCATTAAGGACAGTCTG -3' 5'- TTAGAACTTCAGACCAGCCTGGAC -3'
-----	--

Strain	Genotype	Available from
5905	w[1118]	BDSC
JMT49	w[1118]; sad[10D104], st, e/TM3, P{w[+mC]=GAL4-twi.G}2.3, P{UAS-2xEGFP}AH2.3, Sb[1] Ser[1]	Tennessee Lab
JMT54	w[1118]; sad [5F45] ru, h, th, st, cu, sr, e/TM3, P{w[+mC]=GAL4-twi.G}2.3, P{UAS-2xEGFP}AH2.3, Sb[1] Ser[1]	Tennessee Lab

Acknowledgments: I thank Dr. Michael B. O'Connor for providing strains containing the sad alleles, the Bloomington Drosophila Stock Center for fly stocks (NIH P40OD018537), the Drosophila Genomics Resource Center (NIH 2P40OD010949) for the cDNA construct used in our analysis, and Flybase (NIH U41HG000739).

References

- Abu-Shumays RL, Fristrom JW. 1997. IMP-L3, A 20-hydroxyecdysone-responsive gene encodes *Drosophila* lactate dehydrogenase: structural characterization and developmental studies. *Dev Genet* 20: 11-22. PMID: 9094207.
- Chávez VM, Marqués G, Delbecque JP, Kobayashi K, Hollingsworth M, Burr J, Natzle JE, O'Connor MB. 2000. The *Drosophila* disembodied gene controls late embryonic morphogenesis and codes for a cytochrome P450 enzyme that regulates embryonic ecdysone levels. *Development* 127: 4115-26. PMID: 10976044.
- Currie PD, Sullivan DT. 1994. Structure and expression of the gene encoding phosphofructokinase (PFK) in *Drosophila melanogaster*. *J Biol Chem* 269: 24679-87. PMID: 7929140.
- Currie PD, Sullivan DT. 1994. Structure, expression and duplication of genes which encode phosphoglyceromutase of *Drosophila melanogaster*. *Genetics* 138: 352-63. PMID: 7828819.
- Gillette CM, Tennessen JM, Reis T. 2021. Balancing energy expenditure and storage with growth and biosynthesis during *Drosophila* development. *Dev Biol* 475: 234-244. PMID: 33582116.
- Karim FD, Thummel CS. 1992. Temporal coordination of regulatory gene expression by the steroid hormone ecdysone. *EMBO J* 11: 4083-93. PMID: 1382981.
- Kovalenko EV, Mazina MY, Krasnov AN, Vorobyeva NE. 2019. The *Drosophila* nuclear receptors EcR and ERR jointly regulate the expression of genes involved in carbohydrate metabolism. *Insect Biochem Mol Biol* 112: 103184. PMID: 31295549.
- Kozlova T, Thummel CS. 2003. Essential roles for ecdysone signaling during *Drosophila* mid-embryonic development. *Science* 301: 1911-4. PMID: 12958367.
- Li H, Tennessen JM. 2017. Methods for studying the metabolic basis of *Drosophila* development. *Wiley Interdiscip Rev Dev Biol* 6: 10.1002/wdev.280. PMID: 28556549.
- Madhavan K, Fox DJ, Ursprung H. 1972. Developmental genetics of hexokinase isozymes in *Drosophila melanogaster*. *J Insect Physiol* 18: 1523-30. PMID: 4626548.
- Maróy P, Kaufmann G, Dübendorfer A. 1988. Embryonic ecdysteroids of *Drosophila melanogaster*. *J Insect Physiol* 34: 633-637. DOI: [https://doi.org/10.1016/0022-1910\(88\)90071-6](https://doi.org/10.1016/0022-1910(88)90071-6)
- Ostberg T, Jacobsson M, Attersand A, Mata de Urquiza A, Jendeberg L. 2003. A triple mutant of the *Drosophila* ERR confers ligand-induced suppression of activity. *Biochemistry* 42: 6427-35. PMID: 12767224.
- Roselli-Rehffuss L, Ye F, Lissemore JL, Sullivan DT. 1992. Structure and expression of the phosphoglycerate kinase (*Pgk*) gene of *Drosophila melanogaster*. *Mol Gen Genet* 235: 213-20. PMID: 1465095.
- Ruud AF, Lam G, Thummel CS. 2010. The *Drosophila* nuclear receptors DHR3 and betaFTZ-F1 control overlapping developmental responses in late embryos. *Development* 137: 123-31. PMID: 20023167.

11/30/2021 - Open Access

- Shaw-Lee R, Lissemore JL, Sullivan DT, Tolan DR. 1992. Alternative splicing of fructose 1,6-bisphosphate aldolase transcripts in *Drosophila melanogaster* predicts three isozymes. *J Biol Chem* 267: 3959-67. PMID: 1740444.
- Shaw-Lee RL, Lissemore JL, Sullivan DT. 1991. Structure and expression of the triose phosphate isomerase (*Tpi*) gene of *Drosophila melanogaster*. *Mol Gen Genet* 230: 225-9. PMID: 1720860.
- Sullivan AA, Thummel CS. 2003. Temporal profiles of nuclear receptor gene expression reveal coordinate transcriptional responses during *Drosophila* development. *Mol Endocrinol* 17: 2125-37. PMID: 12881508.
- Sun XH, Tso JY, Lis J, Wu R. 1988. Differential regulation of the two glyceraldehyde-3-phosphate dehydrogenase genes during *Drosophila* development. *Mol Cell Biol* 8: 5200-5. PMID: 3149711.
- Tennessen JM, Bertagnolli NM, Evans J, Sieber MH, Cox J, Thummel CS. 2014. Coordinated metabolic transitions during *Drosophila* embryogenesis and the onset of aerobic glycolysis. *G3 (Bethesda)* 4: 839-50. PMID: 24622332.
- Tennessen JM, Baker KD, Lam G, Evans J, Thummel CS. 2011. The *Drosophila* estrogen-related receptor directs a metabolic switch that supports developmental growth. *Cell Metab* 13: 139-48. PMID: 21284981.
- Tixier V, Bataillé L, Etard C, Jagla T, Weger M, Daponte JP, Strähle U, Dickmeis T, Jagla K. 2013. Glycolysis supports embryonic muscle growth by promoting myoblast fusion. *Proc Natl Acad Sci U S A* 110: 18982-7. PMID: 24191061.
- Warren JT, Petryk A, Marques G, Jarcho M, Parvy JP, Dauphin-Villemant C, O'Connor MB, Gilbert LI. 2002. Molecular and biochemical characterization of two P450 enzymes in the ecdysteroidogenic pathway of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* 99: 11043-8. PMID: 12177427.
- White KP, Rifkin SA, Hurban P, Hogness DS. 1999. Microarray analysis of *Drosophila* development during metamorphosis. *Science* 286: 2179-84. PMID: 10591654.
- Wright DA, Shaw CR. 1970. Time of expression of genes controlling specific enzymes in *Drosophila* embryos. *Biochem Genet* 4: 385-94. PMID: 5477232.

Funding: J.M.T. is supported by a R35 MIRA award from the National Institute of General Medical Sciences of the National Institutes of Health (R35GM119557).

Author Contributions: Jason M. Tennessen: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Validation, Visualization, Writing - original draft, Writing - review and editing.

Reviewed By: Anonymous

History: Received September 20, 2021 **Revision received** November 14, 2021 **Accepted** November 15, 2021 **Published** November 30, 2021

Copyright: © 2021 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Tennessen, JM (2021). Ecdysone and 20-hydroxyecdysone are not required to activate glycolytic gene expression in *Drosophila melanogaster* embryos. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000501>