


Loss of *eif-2alpha* phosphorylation on S49 (mammalian S51) associated with the integrated stress response hastens development in *C. elegans*

Jarod Rollins^{1§}, Noah Lind¹ and Aric N. Rogers¹¹Mount Desert Island Biological Laboratory, 159 Old Bar Harbor Rd Salisbury Cove, ME 04672, United States of America.

§To whom correspondence should be addressed: jrollins@mdibl.org



Description

The gene *eif-2alpha* (Y37E3.10) in *C. elegans* encodes for the alpha subunit of eukaryotic translation initiation factor 2 (eIF2). The eIF2 trimer is required for delivering charged tRNA^{Met} to the ribosome in a reaction requiring the hydrolysis of GTP to GDP (Sokabe et al., 2012). The phosphorylation of *eif2-alpha* on its 49th serine residue (Nukazuka et al., 2008) prevents replenishing GDP with GTP and thus reduces translation initiation by preventing delivery of tRNA^{Met}. The phosphorylation of the mammalian ortholog of *eif-2alpha* occurs on serine 51 in response to a variety of stresses (Taniuchi et al., 2016) including: osmotic stress, ultraviolet light, cold shock, oxidative stress, heat shock, anoxia, and serum starvation.

In *C. elegans*, the phosphorylation of *eif-2alpha* has shown to increase in response to ER stress (Howard et al., 2016; Richardson et al., 2011), osmotic stress (Lee and Strange, 2012), and uncharged tRNAs (Rousakis et al., 2013). This is thought to promote survival by reducing translation in a way that also promotes longevity. Using CRISPR-cas9 genome engineering (Paix et al., 2015) we replaced serine 49 in *eif-2alpha* with alanine, which cannot be phosphorylated. The *C. elegans* line ANR165 harbors this engineered allele in the wild-type N2 background.

The phosphorylation status of *eif-2alpha* in response to hypertonicity, heat, and oxidative stress was probed using an antibody specific to its phosphorylated form. Wild type N2 worms showed an increased in phosphorylated *eif2-alpha* in response to all stresses tested (Figure 1A, dotted box outlines P-*eif2-alpha* bands). Total protein is shown as a loading control. In comparison, phosphorylated *eif-2alpha* was not detectable in ANR165 treated with any of the stresses. This result confirms that serine 49 is the site of phosphorylation in *eif-2alpha* in response to salt stress and is a novel finding for heat and oxidative stress in *C. elegans*.

In addition, *eif2-alpha* S49A mutants were viable, demonstrating that loss of *eif-2alpha* serine 49 phosphorylation is non-lethal. Furthermore, time to develop from egg to egg laying adult was significantly (Wilcoxon rank sum test, $p < 0.01$, $n=15$) shortened in *eif2-alpha* S49A mutants (Figure 1B) by a mean of 4 hours compared to N2. This result suggests that the phosphorylation status of *eif2-alpha* serves a role in development which may be related to its effect on translation. A role for *eif2-alpha* in development is not surprising as mutants expressing a phosphomimetic version of it in the ASI neurons were impaired in growth and development (Kulalert et al., 2017).

Methods

[Request a detailed protocol](#)

C. elegans strains were maintained at 20°C on NMG plates spotted with OP50 as a food source. Hypertonic stress was induced by placing worms on NGM plates containing 200 mM sodium chloride for 3 h. Heat stress was induced by placing worms at 35°C for 1 h. Oxidative stress was induced by placing worms on NMG plates containing 120 μM juglone for 3h. All stresses were performed in the presence of OP50 on day 1 adult worms synchronized by timed egg lays. 10 μg of total protein as determined from DC protein assay (Bio-Rad) from each sample was separated on 4-20%

11/22/2017 - Open Access

mini-Protean TGX stain-free gels (Bio-Rad). The resulting gels were exposed to UV for 5 m to allow fluorescent detection of total protein present.

Reagents

Strains: N2, ANR165 : *eif-2alpha(rog3[S49A])*

Antibody: Phospho-eif2 α (Ser51) #9721, Cell Signaling Technology

References

Howard AC, Rollins J, Snow S, Castor S, Rogers AN. Reducing translation through eIF4G/IFG-1 improves survival under ER stress that depends on heat shock factor HSF-1 in *Caenorhabditis elegans*. *Aging Cell*. 2016;15:1027–38. DOI: 10.1111/accel.12516 | PMID: 27538368.

Kulalert W, Sadeeshkumar H, Zhang YK, Schroeder FC, Kim DH. Molecular Determinants of the Regulation of Development and Metabolism by Neuronal eIF2 α Phosphorylation in *Caenorhabditis elegans*. *Genetics*. 2017;206:251–63. DOI: 10.1534/genetics.117.200568 | PMID: 28292919.

Lee EC-H, Strange K. GCN-2 dependent inhibition of protein synthesis activates osmosensitive gene transcription via WNK and Ste20 kinase signaling. *AJP Cell Physiol*. 2012;303:C1269–77. DOI: 10.1152/ajpcell.00294.2012 | PMID: 23076791.

Nukazuka A, Fujisawa H, Inada T, Oda Y, Takagi S. Semaphorin controls epidermal morphogenesis by stimulating mRNA translation via eIF2 α in *Caenorhabditis elegans*. *Genes Dev*. 2008;22:1025–36. DOI: 10.1101/gad.1644008 | PMID: 18413715.

Paix A, Folkmann A, Rasoloson D, Seydoux G. High Efficiency, Homology-Directed Genome Editing in *Caenorhabditis elegans* Using CRISPR/Cas9 Ribonucleoprotein Complexes. *Genetics*. 2015. DOI: 10.1534/genetics.115.179382 | PMID: 26187122.

Richardson CE, Kinkel S, Kim DH. Physiological IRE-1-XBP-1 and PEK-1 Signaling in *Caenorhabditis elegans* Larval Development and Immunity. 2011;7:e1002391. DOI: 10.1371/journal.pgen.1002391 | PMID: 22125500.

Rousakis A, Vlassis A, Vlanti A, Patera S, Thireos G, Syntichaki P. The general control nonderepressible-2 kinase mediates stress response and longevity induced by target of rapamycin inactivation in *Caenorhabditis elegans*. *Aging Cell*. 2013;12:742–51. DOI: 10.1111/accel.12101 | PMID: 23692540.

Sokabe M, Fraser CS, Hershey JWB. The human translation initiation multi-factor complex promotes methionyl-tRNAi binding to the 40S ribosomal subunit. *Nucleic Acids Res*. 2012;40:905–13. DOI: 10.1093/nar/gkr772 | PMID: 21940399.

Taniuchi S, Miyake M, Tsugawa K, Oyadomari M, Oyadomari S. Integrated stress response of vertebrates is regulated by four eIF2 α kinases. *Sci Rep*. 2016;6:srep32886. DOI: 10.1038/srep32886 | PMID: 27633668.

Funding: This work was supported by grants from the National Institute on Aging of the National Institutes of Health (R21AG056743) and by the Ellison Medical Foundation (AG-NS-1087-13). This research was also supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant numbers P20GM103423 and P20GM104318.

Reviewed By: Elaine Lee

History: Received October 20, 2017 Accepted November 16, 2017 Published November 22, 2017

Copyright: © 2017 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Rollins, J; Lind, N; Rogers, AN (2017). Loss of *eif-2alpha* phosphorylation on S49 (mammalian S51) associated with the integrated stress response hastens development in *C. elegans*. *microPublication Biology*. <https://doi.org/10.17912/W2BM1S>