

# Toxicity of secondary metabolites of *Paenarthrobacter nicotinovorans*

Justyna Kakol<sup>1</sup>, Mainor Vang<sup>1</sup>, Drew Sausen<sup>1</sup>, Tracey Steeno<sup>1</sup>, Angelo Kolokithas<sup>1§</sup> <sup>1</sup>Northeast Wisconsin Technical College, Green Bay, Wisconsin, United States <sup>§</sup>To whom correspondence should be addressed: angelo.kolokithas@nwtc.edu

## Abstract

In a previous study, *Paenarthrobacter nicotinovorans* was isolated and screened for antimicrobial activity. Many promising antibiotic candidates are not ultimately used because of toxicity. After screening secondary metabolites for antimicrobial activity, they were screened for cytotoxicity *in vitro* in the human HeLa cell line. The results show that there is no detectable cytotoxicity in HeLa cells.



Figure 1. Toxicity of Paenarthrobacter nicotinovorans secondary metabolites on HeLa cells:

Relative fluorescence units (RFU) measured for the following samples: positive control (digitonin), solvent control (methanol), negative control (water), and three different concentrations of the secondary metabolite (10  $\mu$ L, 20  $\mu$ L, and 30  $\mu$ L). The error bars represent standard error from the mean and the program used to calculate p values was Microsoft Excel using a t-test. The p values concerning the difference between digitonin and water was 6.92 x 10<sup>-6</sup>, digitonin and methanol was 3.61 x 10<sup>-6</sup>, digitonin and 10 $\mu$ l of metabolite was 3.34 x 10<sup>-6</sup>, digitonin and 20 $\mu$ l of metabolite was 7.60 x 10<sup>-6</sup>, and digitonin and 30 $\mu$ l of metabolite was 6.48 x 10<sup>-6</sup>.

### Description

Upon finding that the secondary metabolites of *Paenarthrobacter nicotinovorans* were displaying antimicrobial activity (Kakol et al., 2023), the toxicity of the metabolites were tested *in vitro*. The CytoTox-Fluor<sup>TM</sup> Cytotoxicity Assay was utilized to test the cytotoxicity of increasing amounts of the secondary metabolites as compared to the toxicity of digitonin, methanol, and water (Figure 1). Increasing the amount of secondary metabolites did not result in increased toxicity as compared to the digitonin control and were more like the negative controls.

Though *Paenarthrobacter nicotinovorans* is described in the literature being involved in nicotine metabolism, its antimicrobial activity has only been tested by the production of silver nanoparticles. (Huq and Akter, 2021) One limitation of this experiment is only one measure of cytotoxicity was utilized, though there are many more available on the market to identify different types of toxicity. Further studies should utilize these measures to identify toxicity that this assay may have missed. Nonetheless, *Paenarthrobacter nicotinovorans* represents a good candidate for further antimicrobial development as it seems not to display toxicity in Hela cells.

## microPublication BIOLOGY

#### 9/19/2023 - Open Access

#### **Methods**

#### Cell culture

Hela cells (ATCC #CCL-2) were maintained in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1x Penicillin-Streptomycin-Neomycin (PSN).

#### Toxicity testing

The toxicity of the secondary metabolites from the *Paenarthrobacter nicotinovorans* bacteria to the HeLa cells was measured by using a commercial CytoTox-Fluor<sup>TM</sup> Cytotoxicity Assay (Promega Catalog #G9260) according to the manufacturer's instructions. The positive control was 300ug/ml Digitonin (Orczyk et al, 2017) (Invitrogen, BN2006). Methanol (Sigma, 67-56-1) and DI H2O were used as vehicle and negative controls, respectively. The extract of *Paenarthrobacter nicotinovorans* (10mg/ml) was used at various concentrations in the toxicity experiments, and various incubation times were used to determine if concentration or incubation time increased toxicity. Ten thousand Hela Cells per well were used in a 96-well plate and the fluorescence measurements were obtained using the Varioskan<sup>TM</sup> LUX (ThermoFischer) multimode microplate reader with the rhodamine-110 (485nmEx/520Em) filter set.

#### Statistics

Microsoft Excel was used to analyze the data. Student T-tests were utilized to determine the level of significance (p values). All experiments and samples were run in triplicate.

**Acknowledgements:** The authors would like to thank Northeast Wisconsin Technical College for supporting this initiative at the college. The Tiny Earth organization for the protocols and tester strains of bacteria, Drs. Matthew Petersen and Troy Moore for their thoughtful review of this manuscript.

#### References

Huq MA, Akter S. 2021. Bacterial Mediated Rapid and Facile Synthesis of Silver Nanoparticles and Their Antimicrobial Efficacy against Pathogenic Microorganisms. Materials (Basel) 14(10). PubMed ID: <u>34069757</u>

Kakol J, Vang M, Sausen D, Steeno T, Kolokithas A. 2023. Antimicrobial activity of Paenarthrobacter nicotinovorans. MicroPubl Biol 2023. PubMed ID: <u>37705710</u>

Orczyk M, Wojciechowski K, Brezesinski G. 2017. Disordering Effects of Digitonin on Phospholipid Monolayers. Langmuir 33(15): 3871-3881. PubMed ID: <u>28333465</u>

#### Funding: No award

**Author Contributions:** Justyna Kakol: investigation, methodology, visualization, data curation, conceptualization. Mainor Vang: data curation, conceptualization, investigation, methodology. Drew Sausen: methodology, resources, project administration, data curation. Tracey Steeno: resources, project administration, validation. Angelo Kolokithas: conceptualization, data curation, formal analysis, investigation, methodology, project administration, resources, supervision, validation, writing - original draft, writing - review editing.

#### Reviewed By: Kurt Regner, Anonymous

History: Received July 13, 2023 Revision Received August 24, 2023 Accepted September 18, 2023 Published Online September 19, 2023 Indexed October 3, 2023

**Copyright:** © 2023 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Kakol, J; Vang, M; Sausen, D; Steeno, T; Kolokithas, A (2023). Toxicity of secondary metabolites of *Paenarthrobacter nicotinovorans*. microPublication Biology. <u>10.17912/micropub.biology.000923</u>