

Xrn1 biochemically associates with eisosome proteins after the post diauxic shift in yeast

Baptiste Courtin¹, Abdelkader Namane¹, Maite Gomard¹, Laura Meyer¹, Alain Jacquier¹, Micheline Fromont-Racine^{1§}

¹Institut Pasteur, Cytoplasmic mRNA surveillance in yeast, Centre National de la Recherche Scientifique, UMR 3525, 75724 Paris Cedex 15, France

[§]To whom correspondence should be addressed: mfromont@pasteur.fr

Abstract

mRNA degradation is one of the main steps of gene expression, and a key player is the 5'-3' exonuclease Xrn1. In *Saccharomyces cerevisiae*, it was previously shown, by a microscopy approach, that Xrn1 is located to different cellular compartments, depending on physiological state. During exponential growth, Xrn1 is distributed in the cytoplasm, while it co-localizes with eisosomes after the post-diauxic shift (PDS). Here, we biochemically characterize the Xrn1-associated complexes in different cellular states. We demonstrate that, after PDS, Xrn1 but not the decapping nor Lsm1-7/Pat1 complexes associates with eisosomal proteins, strengthening the model that sequestration of Xrn1 in eisosomes preserves mRNAs from degradation during PDS.




Figure 1. :

Composition of Xrn1-TAP and Pil1-TAP complexes in different conditions. **A.** Affinity purified Xrn1-TAP associated complexes were subjected to LC-MS/MS identification and quantification. Each panel compares Xrn1-TAP eluate (y-axis) with the total cellular extract (input) (x-axis) from a cell culture in different physiological conditions, as indicated. Each grey dot indicates an identified protein. Red and yellow dots indicate eisosome proteins, and Lsm1-7/Pat1 proteins, respectively, and purple triangles represent the DCP proteins. Proteins of the large (RPLs) or the small ribosomal subunit (RPSs) are indicated by blue and green dots, respectively. The black diamond designates the Xrn1-TAP bait. **B.** Affinity purification using Xrn1-TAP as bait in PDS condition. Volcano plot shows the fold change (\log_2 LFQ cpx/LFQ input) on the x-axis and the p -value distribution ($-\log_{10} p$ -value) on the y-axis for the proteins identified in the affinity purifications. Colored symbols are as indicated in A. **C.** Comparison of affinity purifications using Xrn1-TAP as bait in exponential and PDS conditions; symbols as in A. **D.** Affinity purifications using Pil1-TAP as bait in PDS condition, as in B. **E.** Scatter plot which compares Xrn1-TAP eluate with the cellular lysate (input) in the absence of Pil1 in PDS; symbols as in A. Purifications for the panel B, C and D were done in triplicate, and in duplicate for the panel E.

Description

Gene expression is a crucial and well-orchestrated interplay between transcription, translation, and mRNAs and proteins decay. In eukaryotes, cytoplasmic mRNA decay is intimately linked to translation and mediated for a large part by the 5'-3' mRNA degradation pathway. After mRNA decapping by the DCP complex, with the assistance of its co-factors, the mRNA is degraded by the exonuclease Xrn1 (Parker 2012; Fromont-Racine and Saveanu 2014; for review). This enzyme is highly processive and requires a 5' monophosphorylated RNA end, produced by decapping (Stevens 1988; Larimer and Stevens 1990). The coupling between mRNA degradation and translation is now well established. First, analyzes of yeast cellular extracts separated on sucrose gradient suggested that the 5' mRNA degradation could occurs co-translationally (Hu et al. 2009). Second, mapping of the 5'-end by RNA-Seq revealed that degradation intermediates are phased according to a typical codon-specific trinucleotide shift pattern. This observation strongly indicated that at least a fraction of mRNAs degradation occurred during translation (Pelechano et al. 2015). Finally, a cryo-electron microscopy structure of yeast 80S ribosome-Xrn1 nuclease complex showed that Xrn1 can be intimately and specifically bound to the mRNA exit tunnel region of the ribosome, fully consistent with the fact that 5'-3' mRNA degradation pathway is physically coupled to translation (Tesina et al. 2019).

Gene expression varies, depending on the physiological state of the cells. The *S. cerevisiae* cells must quickly adapt to survive stress conditions. Consequently, a fast reprogramming of gene expression must be set up (DeRisi et al. 1997). Under nutrient starvation, translation of many mRNAs is inhibited (Ashe et al. 2000). This is correlated with Xrn1 cellular localization changes. In response to stress conditions, such as glucose depletion, many mRNA-associated factors involved in mRNA degradation, such as Xrn1 and the DCP complex, are concentrated in processing bodies (P-bodies) (Teixeira and Parker 2007). During the diauxic shift, which happens through the transition from exponential growth to stationary phase, as well as during the post-diauxic shift period (PDS), many cellular modifications occur. The Malinsky's team previously observed that Xrn1 is sequestered in the eisosome in PDS by a microscopy approach (Grousl et al. 2015; Vaškovičová et al. 2017).

The eisosome is a structure associated with the plasma membrane near endocytosis sites. The two main components, Pil1 and Lsp1, share 72% sequence identity and are present in around equimolar proportions. However, despite this strong identity, Pil1 appears to be the major actor of eisosome biogenesis (Walther et al. 2006; Fröhlich et al. 2014). To be efficient, Pil1 and Lsp1 are phosphorylated by two protein kinases, Pkh1 and Pkh2, which are part of the eisosome (Walther et al. 2007). When these two kinases are mutated, the eisosome structure is impaired and endocytosis sites localization is affected suggesting a link between eisosome and endocytosis (Luo et al. 2008). However, the role of the eisosome is far from being understood.

We first identified proteins associated with Xrn1 complex during different physiological states of the cell. We compared the Xrn1-associated complexes obtained by affinity purifications from exponential state extracts ($OD_{600\text{ nm}} 0.6$ and $OD_{600\text{ nm}} 2$), diauxic shift (DS) and Post-diauxic shift (PDS) extracts (Figure 1A). LC-MS-MS identification followed by a label-free quantification (LFQ) analysis revealed that the eisosome proteins (red dots), were not enriched with Xrn1-complexes in exponential phase but started to be weakly enriched in DS and were highly enriched with the Xrn1-associated complexes from PDS cells. Among the eisosome proteins, the two main proteins of this structure, Pil1 and Lsp1, were the most abundant and the most enriched with the PDS Xrn1-complexes. In contrast, the DCP complex (Figure 1A, purple triangles) was less and less present in the Xrn1-associated complexes, from the exponential phase to the PDS, while the presence of the Lsm1-7/Pat1 complex (Figure 1A, yellow dots) was still enriched with Xrn1 (Figure 1A, Dataset 1). A volcano-plot representation revealed the Xrn1-associated complex composition in PDS (Figure 1B, Dataset 1). It highlighted that the most enriched proteins during PDS are the components of the eisosome (red dots). The Lsm1-7/Pat1 proteins (yellow dots) were also enriched but not those of the DCP complex (purple triangles) nor the ribosome (blue and green dots). A comparison between the Xrn1-TAP associated complexes in exponential phase and PDS (Figure 1C, Dataset 1) distinctly showed that Xrn1 was highly associated

with the eisosome in PDS (left part of the graph) and highly associated with the DCP complex and with the ribosome in exponential phase (right part of the graph).

To biochemically validate the presence of Xrn1 in the eisosome in PDS and to determine if the Lsm1-7/Pat1 complex is also present in the eisosome, we performed an affinity purification using Pil1-TAP as bait in PDS conditions (Figure 1D, Dataset 1). LC-MS-MS identification followed by a quantitative LFQ analysis indicated that Xrn1 (black diamond) was highly enriched with Pil1 while the Lsm1-7/Pat1 complex was not. In addition, we observed that, in the absence of Pil1, Xrn1 was still associated with the Lsm1-7/Pat1 complex, whereas the eisosome components were missing (Figure 1E, Dataset 1). This is consistent with the observation that in *pil1Δ* mutant strain, the eisosome is completely disassembled (Grousl et al. 2015). In conclusion, we confirmed here, by a biochemical approach, that Xrn1 is localized in the eisosome after post diauxic shift, whereas neither the DCP complex nor the Lsm1-7/Pat1 factors are present in this structure. Our results also suggest that, after PDS and independently of the eisosome, a part of Xrn1 remains associated with the Lsm1-7/Pat1 complex, but not with the DCP complex.

Altogether, these results indicate that, during PDS, Xrn1 is still present but mainly sequestered in the eisosome, probably to protect the mRNAs from full degradation. Nevertheless, a fraction of Xrn1 seems to remain associated with the Lsm1-7/Pat1 complex. It would be of interest to investigate the role of this interaction, independent of the eisosome and to determine if the interactions between Xrn1 and the Lsm1-7/Pat1 complex and those between Xrn1 and the eisosome are RNA-mediated. This study suggests that sequestration of Xrn1 plays a role in maintaining cell homeostasis during adaptation to nutrient starvation.

Methods

Yeast strains and culture conditions. All the strains derive from the BY4741 strain (*MATa*, *ura3Δ0*, *his3Δ1*, *leu2Δ0*, *met15Δ0*). LMA5015, Xrn1-TAP:HIS3MX6 and LMA5434, Pil1-TAP:HIS3MX6 are from the TAP collection (Ghaemmaghami et al. 2003). LMA5512, Xrn1-TAP:HIS3MX6 *pil1Δ::KANMX4* was constructed by transformation of the LMA5015 strain with a PCR fragment obtained with *pil1Δ::KANMX4* strain from the Euroscarf collection using oligonucleotide fw: GAATGGACACTAGACTCTGC and oligonucleotide rv: GGGAACAGAAATGATTATCTGTCC. The cell cultures were grown at 30°C in YPGlu medium, up to an OD_{600nm} 0.6 or 2.0. To collect the DS and PDS samples, the cultures were diluted to an OD_{600nm} 0.005 and grown until they reached the Diauxic State (DS) after 19h of culture or Post Diauxic State (PDS) after 37h of culture. 4000 OD_{600nm} of cell cultures were collected, washed with cold water and stored at -80°C.

Affinity purification. The cells were resuspended with lysis buffer (20mM Hepes pH7.4, 10mM MgCl₂, 100mM KOAc) containing a protease-inhibiting reagent (Roche) (1mL per mg of cells). Then, 500 μL of acid-washed glass beads were added per mL of cell suspension and vortexed three-time 40 sec. at 4°C, 6m/sec. (MP FastPrep™, Fisher Scientific). The suspension was centrifuged for 20 min. at 4°C, 4000 rpm. 0.5% Triton was added to the supernatant, followed by 25 μL of covalently coupled IgG-Dynabeads® magnetic beads resuspended in lysis buffer. The lysate was then incubated for 2 hours at 4°C under agitation. The magnetic beads were recovered and washed five times in washing buffer (20mM Hepes pH7.4, 10mM MgCl₂, 100mM KOAc, 0.5% Triton), then once in lysis buffer. They were resuspended in 2% SDS, 1X TE and eluted by incubation at 65°C for 15 min. The SDS was removed with HiPPR™ Detergent Removal Resin kit (Thermo Fisher Scientific). Proteins were precipitated by Methanol/Chloroform method (Wessel and Flügge 1984).

Mass spectrometry analysis. The samples were treated as described in (Defenouillère et al. 2013). Briefly, endoprotease Lys-C and Trypsin digested peptides were analyzed on an LTQ-Orbitrap Velos mass spectrometer (Thermo Fisher Scientific, Bremen). The raw data were analyzed with MaxQuant (version 2.0.3.0). Only proteins identified with at least 2 peptides were selected for further quantification analysis. The abundance of the identified proteins was achieved in the Perseus environment (version 1.6.14). Protein group LFQ intensities were log₂ transformed. After comparison with a control group, using the Student's t test statistic, the results were represented as volcano plots or scatter plots.

Data availability

Raw mass spectrometry data were posted in the ProteomeXchange Consortium via the PRIDE repository. The accession numbers are PXD044514, PXD044631, PXD044634, PXD044653.

Acknowledgements: We are grateful to the proteomics platform of the Institut Pasteur for the availability of the Orbitrap Velos. We thank Gwenaél Badis and Cosmin Saveanu for discussions and criticism on the manuscript. We are grateful to Lucia Oreus for providing media and buffers used in this study and Christelle Lenormand for her administrative assistance. We thank Emmanuelle Permal for uploading the proteomic data in the ProteomeXchange Consortium.

Extended Data

Description: Data from mass spectrometry analysis. Resource Type: Dataset. File: [Dataset1.xlsx](#). DOI: [10.22002/30gmd-r4w62](#)

References

- Ashe, M. P., De Long, S. K., Sachs, A. B. 2000. Glucose depletion rapidly inhibits translation initiation in yeast. *Molecular Biology of the Cell*. 11: 833. DOI: [10.1091/mbc.11.3.833](#)
- Defenouillère, Quentin, Yao, Yanhua, Mouaikel, John, Namane, Abdelkader, Galopier, Aurélie, Decourty, Laurence, et al., Fromont-Racine, Micheline. 2013. Cdc48-associated complex bound to 60S particles is required for the clearance of aberrant translation products. *Proceedings of the National Academy of Sciences of the United States of America*. 110: 5046. DOI: [10.1073/pnas.1221724110](#)
- DeRisi, Joseph L., Iyer, Vishwanath R., Brown, Patrick O. 1997. Exploring the Metabolic and Genetic Control of Gene Expression on a Genomic Scale. *Science*. 278: 680. DOI: [10.1126/science.278.5338.680](#)
- Fröhlich, Florian, Christiano, Romain, Olson, Daniel K., Alcazar-Roman, Abel, DeCamilli, Pietro, Walther, Tobias C. 2014. A role for eisosomes in maintenance of plasma membrane phosphoinositide levels. *Molecular Biology of the Cell*. 25: 2797. DOI: [10.1091/mbc.E13-11-0639](#)
- Fromont-Racine, Micheline, Saveanu, Cosmin 2014. mRNA degradation and decay. *Fungal RNA Biology*. : 159.
- Ghaemmaghami, Sina, Huh, Won-Ki, Bower, Kiowa, Howson, Russell W., Belle, Archana, Dephoure, Noah, O'Shea, Erin K., Weissman, Jonathan S. 2003. Global analysis of protein expression in yeast. *Nature*. 425: 737. DOI: [10.1038/nature02046](#)
- Grousl, Tomas, Opekarová, Miroslava, Stradalova, Vendula, Hasek, Jiri, Malinsky, Jan 2015. Evolutionarily Conserved 5'-3' Exoribonuclease Xrn1 Accumulates at Plasma Membrane-Associated Eisosomes in Post-Diauxic Yeast. *PLoS ONE*. 10: e0122770. DOI: [10.1371/journal.pone.0122770](#)
- Hu, Wenqian, Sweet, Thomas J., Chamnongpol, Sangpen, Baker, Kristian E., Coller, Jeff 2009. Co-translational mRNA decay in *Saccharomyces cerevisiae*. *Nature*. 461: 225. DOI: [10.1038/nature08265](#)
- Larimer, F. W., Stevens, A. 1990. Disruption of the gene XRN1, coding for a 5'----3' exoribonuclease, restricts yeast cell growth. *Gene*. 95: 85. DOI: [10.1016/0378-1119\(90\)90417-p](#)
- Luo, Guangzuo, Gruhler, Albrecht, Liu, Ying, Jensen, Ole N., Dickson, Robert C. 2008. The sphingolipid long-chain base-Pkh1/2-Ypk1/2 signaling pathway regulates eisosome assembly and turnover. *The Journal of Biological Chemistry*. 283: 10433. DOI: [10.1074/jbc.M709972200](#)
- Parker, R. 2012. RNA Degradation in *Saccharomyces cerevisiae*. *Genetics*. 191: 671. DOI: [10.1534/genetics.111.137265](#)
- Pelechano, Vicent, Wei, Wu, Steinmetz, Lars M. 2015. Widespread co-translational RNA decay reveals ribosome dynamics. *Cell*. 161: 1400. DOI: [10.1016/j.cell.2015.05.008](#)
- Stevens, A. 1988. mRNA-decapping enzyme from *Saccharomyces cerevisiae*: purification and unique specificity for long RNA chains. *Molecular and Cellular Biology*. 8: 2005. DOI: [10.1128/mcb.8.5.2005-2010.1988](#)
- Teixeira, Daniela, Parker, Roy 2007. Analysis of P-Body Assembly in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell*. 18: 2274. DOI: [10.1091/mbc.e07-03-0199](#)
- Tesina, Petr, Heckel, Elisabeth, Cheng, Jingdong, Fromont-Racine, Micheline, Buschauer, Robert, Kater, Lukas, et al., Beckmann, Roland. 2019. Structure of the 80S ribosome-Xrn1 nuclease complex. *Nature Structural & Molecular Biology*. 26: 275. DOI: [10.1038/s41594-019-0202-5](#)
- Vaškovičová, Katarína, Awadová, Thuraya, Veselá, Petra, Balážová, Mária, Opekarová, Miroslava, Malinsky, Jan 2017. mRNA decay is regulated via sequestration of the conserved 5'-3' exoribonuclease Xrn1 at eisosome in yeast. *European Journal of Cell Biology*. 96: 591. DOI: [10.1016/j.ejcb.2017.05.001](#)
- Walther, Tobias C, Aguilar, Pablo S, Fröhlich, Florian, Chu, Feixia, Moreira, Karen, Burlingame, Alma L, Walther, Peter 2007. Pkh-kinases control eisosome assembly and organization. *The EMBO Journal*. 26: 4946. DOI: [10.1038/sj.emboj.7601933](#)
- Walther, Tobias C., Brickner, Jason H., Aguilar, Pablo S., Bernales, Sebastián, Pantoja, Carlos, Walther, Peter 2006. Eisosomes mark static sites of endocytosis. *Nature*. 439: 998. DOI: [10.1038/nature04472](#)
- Wessel, D., Flüggé, U. I. 1984. A method for the quantitative recovery of protein in dilute solution in the presence of detergents and lipids. *Analytical Biochemistry*. 138: 141.

9/7/2023 - Open Access

Funding: This work was supported by the ANR-17-CE11-0049-01, the ANR-17-CE12-0024-02 and the ANR-18-CE11-0003-04 Grants from the Agence Nationale de la Recherche, the Institut Pasteur, and the Centre National de la Recherche Scientifique.

Author Contributions: Baptiste Courtin: investigation, formal analysis. Abdelkader Namane: investigation, formal analysis. Maite Gomard: investigation. Laura Meyer: investigation. Alain Jacquier: funding acquisition, writing - review editing. Micheline Fromont-Racine: conceptualization, funding acquisition, supervision, writing - review editing.

Reviewed By: GUILLAUME CHANFREAU

History: **Received** July 19, 2023 **Revision Received** August 29, 2023 **Accepted** August 31, 2023 **Published Online** September 7, 2023 **Indexed** September 21, 2023

Copyright: © 2023 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Courtin, B; Namane, A; Gomard, M; Meyer, L; Jacquier, A; Fromont-Racine, M (2023). Xrn1 biochemically associates with eisosome proteins after the post diauxic shift in yeast. microPublication Biology. [10.17912/micropub.biology.000926](https://doi.org/10.17912/micropub.biology.000926)