

Atypical soluble guanylyl cyclases control brain size in *Drosophila*

Daniel Prieto^{1,2§}, Boris Egger³, Rafael Cantera¹

¹Departamento de Biología del Neurodesarrollo, Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Uruguay

²Departamento de Neurofisiología Celular y Molecular, Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Uruguay

³Department of Biology, University of Fribourg, Fribourg, Switzerland

[§]To whom correspondence should be addressed: dprieto@fcien.edu.uy

Abstract

Hypoxia-induced proliferation of neural stem cells has a crucial role in brain development. In the brain of *Drosophila melanogaster*, the optic lobe exhibits progressive hypoxia during larval development. Here, we investigate an alternative oxygen-sensing mechanism within this brain compartment, distinct from the canonical hypoxia signaling pathway mediated by HIF. Using genetic tools, immunostaining, and confocal microscopy, we demonstrate that the loss of the atypical soluble guanylyl cyclase (asGC) subunit *Gyc88E*, or the ectopic expression of *Gyc89Db* in neural stem cells leads to increased optic lobe volume. We propose the existence of a link between cGMP signaling and neurogenesis in the developing brain.




Figure 1. Effect of atypical soluble guanylyl cyclase loss- and gain-of-function in brain development of *Drosophila* larvae:

(A) Volume of brain hemispheres at third larval instar. Loss-of-function *Gyc88E*^{-/-} mutant larvae (88E^{-/-}) displayed increased brain hemisphere volume, (median±SD: 1.93e+06 ± 3.54e+05 μm³), whereas *Gyc89Da*^{-/-}*Db*^{-/-} double mutant (89Da^{-/-}Db^{-/-}) did not (median±SD: 7.29e+05 ± 4.34e+05 μm³). Increased hemisphere volume was also observed in larvae ectopically expressing *Gyc89Db* in optic-lobe neuroepithelial cells under control of *GAL4*^{c855a} (c855a>89Db, median±SD: 1.84e+06 ± 1.53e+05 μm³) or overexpressing it in neuroblasts under control of *insc-Gal4* (*insc*>89Db, median±SD: 1.88e+06 ± 1.85e+05 μm³), compared to *w*¹¹¹⁸ (median±SD: 9.86e+05 ± 5.70e+05 μm³). **(B)** Volume of central brain. Loss-of-function mutant larvae displayed increased central brain volume, *Gyc88E*^{-/-} (median±SD: 1.08e+06 ± 2.46e+05 μm³), *Gyc89Da*^{-/-}*Db*^{-/-} (median±SD: 8.53e+05 ± 1.81e+05 μm³) as did ectopic expression lines driving *Gyc89Db* in optic-lobe neuroepithelial cells (median±SD: 8.53e+05 ± 6.18e+04 μm³). Overexpression in neuroblasts did not alter volume occupied by neuroblasts (median±SD: 8.41e+05 ± 1.37e+05 μm³), compared to *w*¹¹¹⁸ (median±SD: 5.05e+05 ± 1.70e+05 μm³). **(C)** Volume of the optic lobe (OL). *Gyc88E*^{-/-} mutant larvae did not show increased OL volume (median±SD: 8.04e+05 ± 2.01e+05 μm³), nor did *Gyc89Da*^{-/-}*Db*^{-/-} larvae (median±SD: 6.84e+05 ± 2.46e+05 μm³). Larvae ectopically expressing *Gyc89Db* in neuroepithelial cells showed increased OL volume (median±SD: 1.04e+06 ± 4.21e+05 μm³) whereas those overexpressing *Gyc89Db* in neuroblasts did not (median±SD: 8.72e+05 ± 1.95e+05 μm³) when compared to *w*¹¹¹⁸ (median±SD: 4.81e+05 ± 4.23e+05). **(D)** Assessment of neuroblast number using anti-Deadpan immunostaining. Optic lobe neuroblasts were counted in the brain of larvae from each experimental group. The number of neuroblasts was increased in *Gyc88E*^{-/-} (median±SD: 230.00 ± 67.56) and in *Gyc89Da*^{-/-}*Db*^{-/-} mutant larvae (median±SD: 211.50 ± 56.10). Larvae expressing *Gyc89Db* in neuroepithelial cells exhibited a 5.8-fold increase in their neuroblast number (median±SD: 425.50 ± 39.99). Overexpression in neuroblasts did not alter their number (median±SD: 85.00 ± 26.06) compared to *w*¹¹¹⁸ (median±SD: 73.00 ± 79.18) larvae. **(E)** Quantification of proliferating optic lobe neuroblasts, identified by double staining for anti-Deadpan and anti-PH3. The number of proliferating neuroblasts was low, without statistically significant differences in *Gyc88E*^{-/-} (median±SD: 1.00 ± 2.76) or *Gyc89Da*^{-/-}*Db*^{-/-} double mutant larvae (median±SD: 5.00e-01 ± 8.16e-01) compared to *w*¹¹¹⁸ (median±SD: 1.00 ± 7.63). Brains expressing *Gyc89Db* under the neuroepithelial driver *GAL4*^{c855a} (median±SD: 17.5 ± 3.32) showed a 2.5-fold increase in the number of mitotically active neuroblasts, while those expressing *Gyc89Db* under the neuroblast driver *insc-gal4* showed a 3.4-fold increase (median±SD: 24.00 ± 16.76). **(F)** Quantification of total proliferating cells, measured by anti-PH3 immunostaining. *Gyc88E*^{-/-} larvae showed increased numbers of total proliferating cells (1.4-fold, median±SD: 154.00 ± 74.42) whereas *Gyc89Da*^{-/-}*Db*^{-/-} did not (median±SD: 79.50 ± 74.46). Larvae expressing *Gyc89Db* in neuroepithelial cells under control of the *GAL4*^{c855a} driver exhibited a 4.3-fold increase in the number of mitotic cells (median±SD: 459.50 ± 44.14), whilst larvae expressing *Gyc89Db* under the neuroblast driver *insc-gal4* showed a 1.5-fold increase (median±SD: 164.50 ± 65.58) compared to *w*¹¹¹⁸ larvae (median±SD: 64.50 ± 51.07). Mann-Whitney U test, * p<0.05, ** p<0.01, *** p<0.001. Mutant, ectopic expression, and overexpression N=6, *w*¹¹¹⁸ N=12.

Description

Oxygen availability is a powerful driver of evolutionary novelty, and metazoans have developed sophisticated hypoxia sensing systems that are intricately related to the control of stem cell niches and development (Simon and Keith 2008; Rytönen et al. 2011; Hammarlund et al. 2018). Increased atmospheric oxygen levels during evolution had profound effects on insects enabling them, for instance, to develop larger body size and the ability to fly (Stamati et al. 2011).

The atypical soluble guanylyl cyclase (asGC) subunits *Gyc88E*, *Gyc89Da* and *Gyc89Db* are expressed from embryo to adult stages in *Drosophila melanogaster* (Langlais et al. 2004; Morton et al. 2005). They are regulated by O₂ levels and when activated by hypoxia generate cGMP, functioning as molecular O₂ sensors (Vermehren et al. 2006) with faster responses than the canonical hypoxia pathway (de Lima et al. 2021).

The neural stem cells within the optic lobe (OL) of the *Drosophila* larval brain are hypoxic relative to the central brain (Misra et al. 2017), with different cell types experiencing varying degrees of hypoxia (Baccino-Calace et al. 2020). There is evidence that optic lobe progenitor cells might not exhibit a canonical hypoxia response (Baccino-Calace et al. 2020). *In situ* hybridization data (Langlais et al. 2004) suggests that asGC subunits are expressed within the OL, potentially linking O₂ sensing to an alternative signaling pathway activated by hypoxia. Furthermore, transcriptomic data analysis (Yang et al. 2015),

revealed that *Gyc89Db* is expressed in L3 neuroblasts. The combination of different asGC subunits might provide different sensitivity thresholds to hypoxia to *Drosophila* neurons (Lu et al. 2024).

Here, we used genetic tools to investigate the role of asGC in larval brain development. We explored asGC loss-of-function using a *Gyc88E*^{-/-} mutant and a *Gyc89Da*^{-/-}*Db*^{-/-} double mutant. Additionally, we performed targeted gain-of-function with drivers specific to ectopically express *Gyc89Db* in neuroepithelial cells or overexpressing it in neuroblasts.

As the hypoxic OL remains neurogenic during larval life, we hypothesized that this hypoxia might activate asGC. Given that cGMP derived from nitric oxide signaling can activate mammalian neural stem cell proliferation (Santos et al. 2014), if hypoxia-driven cGMP signaling were to activate neural stem cell proliferation, loss of asGC function would abolish it. Contrary to our expectation, the *Gyc88E*^{-/-} loss-of-function mutant exhibited a 1.9-fold increase in brain hemisphere volume (Figure 1A). Interestingly, the double mutation *Gyc89Da*^{-/-}*Db*^{-/-} had not the same effect. Notably, ectopic expression in NE or overexpression of *Gyc89Db* in NB lead to an increase in overall brain size (Figure 1A). The larval brain comprises two functionally and developmentally different compartments, the central brain (CB) which is formed mostly during embryonic stages and contains the neuronal synapses and the optic lobe (OL), which lack fully differentiated neurons with synapses and remains neurogenic throughout larval stages (Hartenstein et al. 2008; Baccino-Calace et al. 2020).

When we measured and compared CB and OL volumes separately, we observed that the macrocephalous phenotype observed in *Gyc88E*^{-/-} mutants can be explained by a specific effect on the CB. *Gyc88E*^{-/-} loss-of-function mutants showed a 2.1-fold increase in CB volume (Figure 1B), while the OL volume remains unaffected (Figure 1C). The double mutation *Gyc89Da*^{-/-}*Db*^{-/-} also elicited an increase in CB volume but limited to a lesser extent (1.7-fold, Figure 1B).

Ectopic expression of *Gyc89Db* under the control of the OL-specific neuroepithelial driver *GAL4*^{c855a} (Egger et al. 2007) resulted in increased volume of both CB (Figure 1B) and OL (Figure 1C). Overexpression under the pan-neuroblast driver *inscuteable*-*GAL4* reproduced the macrocephalous phenotype only in the CB (Fig. 1B). Provided the OL is the major neurogenic region within the larval brain, experiencing a dramatic volume increase between 24-72 hs after larval hatching (Baccino-Calace et al. 2020), we reasoned that, in the larva, an eventual increase in proliferation would only be observable in this hypoxic brain region.

Aiming to further investigate the cellular basis of the macrocephalous phenotype, we assessed neuroblast number and proliferation within the OL. asGC loss-of-function mutants and ectopic expression of *Gyc89Db* in the OL neuroepithelium increased neuroblast number in this region, while overexpression in neuroblasts did not (Figure 1D).

Despite the increased OL volume, no differences in the number of proliferating neuroblasts were observed in the loss-of-function mutants (Figure 1E), whereas *Gyc89Db* ectopic expression in neuroepithelial cells or overexpression in neuroblasts increased the number of mitotically active neuroblasts by 2.5-fold and 3.4-fold, respectively (Figure 1E).

Furthermore, larvae with *Gyc88E*^{-/-} loss of function or ectopic expression of *Gyc89Db* in the neuroepithelium or overexpression in neuroblasts show increased total cell number in mitosis in the optic lobe (Figure 1F), suggesting a link between oxygen-dependent cGMP signaling and the control of proliferation in neural progenitor cells.

Previous research has shown that *Gyc88E* is active in the absence of additional subunits, while *Gyc89Da* and *Gyc89Db* enhanced the activity of *Gyc88E* when co-expressed, suggesting that these enzymes acted likely as heterodimers (Morton and Vermehren 2007; Morton 2011). It is important to mention that *Gyc89Db* is expressed in the brain during larval development (Langlais et al. 2004) and produces cyclic GMP in response to low oxygen *in vitro* (Morton 2004; Vermehren et al. 2006). Thus, asGCs may fine tune the cellular response promoting proliferation through hypoxia-driven cGMP signaling.

Methods

Fly strains

Flies were raised on cornmeal medium at constant temperature 25°C and under 12:12 h light:darkness cycles as previously described (Ferreiro et al. 2018). Loss-of-function stocks *Gyc88E*^{-/-} (Vermehren-Schmaedick et al. 2010) (Z3-1083, from the Seattle tilling project) and the double mutant *Gyc89Da*^{-/-}*Db*^{-/-} (Vermehren-Schmaedick et al. 2010) (BDSC #93108) were a kind gift of David Morton (Oregon Health & Science University, Oregon, USA). The following driver and responder lines were used: *c855a-GAL4* (BDSC stock #6990), *insc-GAL4* (BDSC stock #8751), *UAS-gyc89Db* (Vermehren-Schmaedick et al. 2010) (BDSC stock #93116) was a kind gift from David Morton (Oregon Health & Science University, Oregon, USA). *w*¹¹¹⁸ was used as control as mutant and transgenic lines were built on a *white* background.

Immunostaining and image acquisition

All experiments were performed on wandering third-instar larvae kept in normoxia conditions. Brains were dissected, fixed and immunostained as previously described (Baccino-Calace et al. 2020). Primary antibodies included a monoclonal mouse anti-Discs large antibody to outline all brain cells and allow for precise identification of the OL boundaries (1:20, DSHB #4F3, Developmental Studies Hybridoma Bank (DSHB), Iowa, USA), guinea pig anti-Deadpan to identify neuroblasts (1:2500, kind gift from Jürgen Knoblich, Eroglu et al. 2014) and rabbit anti-phosphorylated H3 histone antibody as a mitotic marker (anti-PH3, 1:200, Cell Signaling Technologies #9713). Fluorescent conjugated secondary antibodies Alexa488, Alexa568 and Cy5 were used (Thermo-Fisher). DNA was stained with 1 µg/ml Hoechst 33342 (Thermo-Fisher). Images were acquired with a Zeiss LSM800 Airyscan confocal microscope and processed with FIJI (Schindelin et al. 2012). Semi-automated OL segmentation was performed with TrakEM2 (Cardona et al. 2012). Analyses and illustrations were made using R version 4.4.0 on RStudio version 2023.06.01.

Reagents

Stocks: Stocks obtained from the Bloomington Drosophila Stock Center (BDSC) or kindly provided by colleagues as stated under the Methods section.

GAL4^{C855} (*w*[1118]; *P*{*w*[+*mW.hs*]=*GawB*}*C855a*)

insc-GAL4 (*w*[*]; *P*{*w*[+*mW.hs*]=*GawB*}*insc*[*Mz1407*])

UAS-gyc89Db (*w*[*]; *P*{*w*[+*mC*]=*UAS-Gyc89Db.V*}2)

Gyc88E^{-/-} (point mutation V474M)

Gyc89Da^{-/-}*Db*^{-/-} (*w*[*]; *P*Bac{*w*[+*mC*]=*RB*}*Gyc89Da*[e01821] *Mi*{*GFP*[*E.3xP3*]=*ET1*}*Gyc89Db*[*MB03197*])

Acknowledgements:

The authors would like to thank Dr. Maria Jose Ferreiro for her continued valuable help with fly care and fruitful discussions, and the Developmental Studies Hybridoma Bank for antibodies. Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study.

References

- Baccino-Calace M, Prieto D, Cantera R, Egger B. 2020. Compartment and cell-type specific hypoxia responses in the developing *Drosophila* brain. *Biology Open* 9: 10.1242/bio.053629. DOI: [10.1242/bio.053629](https://doi.org/10.1242/bio.053629)
- Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, et al., Douglas. 2012. TrakEM2 Software for Neural Circuit Reconstruction. *PLoS ONE* 7: e38011. DOI: [10.1371/journal.pone.0038011](https://doi.org/10.1371/journal.pone.0038011)
- de Lima TbM, Nery LEM, Maciel FbE, Ngo-Vu H, Kozma MT, Derby CD. 2021. Oxygen sensing in crustaceans: functions and mechanisms. *Journal of Comparative Physiology A* 207: 1-15. DOI: [10.1007/s00359-020-01457-z](https://doi.org/10.1007/s00359-020-01457-z)
- Egger B, Boone J Q, Stevens N R, Brand A H, Doe C Q. 2007. Regulation of spindle orientation and neural stem cell fate in the *Drosophila* optic lobe.. *Neural Development*. 2: 1. DOI: [10.1186/1749-8104-2-1](https://doi.org/10.1186/1749-8104-2-1)
- Eroglu E, Burkard TR, Jiang Y, Saini N, Homem CCF, Reichert H, Knoblich JA. 2014. SWI/SNF Complex Prevents Lineage Reversion and Induces Temporal Patterning in Neural Stem Cells. *Cell* 156: 1259-1273. DOI: [10.1016/j.cell.2014.01.053](https://doi.org/10.1016/j.cell.2014.01.053)
- Ferreiro MaJ, Pérez C, Marchesano M, Ruiz S, Caputi A, Aguilera P, Barrio R, Cantera R. 2018. *Drosophila melanogaster* White Mutant *w*1118 Undergo Retinal Degeneration. *Frontiers in Neuroscience* 11: 10.3389/fnins.2017.00732. DOI: [10.3389/fnins.2017.00732](https://doi.org/10.3389/fnins.2017.00732)
- Hammarlund EU, von Stedingk K, Pählman S. 2018. Refined control of cell stemness allowed animal evolution in the oxic realm. *Nature Ecology & Evolution* 2: 220-228. DOI: [10.1038/s41559-017-0410-5](https://doi.org/10.1038/s41559-017-0410-5)
- Hartenstein V, Spindler S, Pereanu W, Fung S. 2008. The development of the *Drosophila* larval brain. *Adv Exp Med Biol* 628: 1-31. PubMed ID: [18683635](https://pubmed.ncbi.nlm.nih.gov/18683635/)
- Langlais KK, Stewart JA, Morton DB. 2004. Preliminary characterization of two atypical soluble guanylyl cyclases in the central and peripheral nervous system of *Drosophila melanogaster*. *Journal of Experimental Biology* 207: 2323-2338. DOI: [10.1242/jeb.01025](https://doi.org/10.1242/jeb.01025)
- Lu S, Qian CS, Grueber WB. 2024. Mechanisms of gas sensing by internal sensory neurons in *Drosophila* larvae. : 10.1101/2024.01.20.576342. DOI: [10.1101/2024.01.20.576342](https://doi.org/10.1101/2024.01.20.576342)

- Misra T, Baccino-Calace M, Meyenhofer F, Rodriguez-Crespo D, Akarsu H, Armenta-Calderón R, et al., Luschnig. 2016. A genetically encoded biosensor for visualizing hypoxia responses *in vivo*. *Biology Open* : 10.1242/bio.018226. DOI: [10.1242/bio.018226](https://doi.org/10.1242/bio.018226)
- Morton DB. 2004. Atypical Soluble Guanylyl Cyclases in *Drosophila* Can Function as Molecular Oxygen Sensors. *Journal of Biological Chemistry* 279: 50651-50653. DOI: [10.1074/jbc.C400461200](https://doi.org/10.1074/jbc.C400461200)
- Morton David B. 2011. Behavioral responses to hypoxia and hyperoxia in *Drosophila* larvae molecular and neuronal sensors. *Fly*. 5: 119. DOI: [10.4161/fly.5.2.14284](https://doi.org/10.4161/fly.5.2.14284)
- Morton DB, Langlais KK, Stewart JA, Vermehren A. 2005. Comparison of the properties of the five soluble guanylyl cyclase subunits in *Drosophila melanogaster*. *Journal of Insect Science* 5: 1-10. DOI: [10.1673/031.005.1201](https://doi.org/10.1673/031.005.1201)
- Morton David B., Vermehren Anke. 2007. Soluble guanylyl cyclases in invertebrates: Targets for NO and O₂. *Advances in Experimental Biology*. 1: 65. DOI: [10.1016/S1872-2423\(07\)01003-4](https://doi.org/10.1016/S1872-2423(07)01003-4)
- Rytönen KT, Storz JF. 2010. Evolutionary origins of oxygen sensing in animals. *EMBO reports* 12: 3-4. DOI: [10.1038/embor.2010.192](https://doi.org/10.1038/embor.2010.192)
- Santos AI, Carreira BP, Nobre RJ, Carvalho CM, Araújo IsM. 2014. Stimulation of Neural Stem Cell Proliferation by Inhibition of Phosphodiesterase 5. *Stem Cells International* 2014: 1-13. DOI: [10.1155/2014/878397](https://doi.org/10.1155/2014/878397)
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al., Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods* 9: 676-682. DOI: [10.1038/nmeth.2019](https://doi.org/10.1038/nmeth.2019)
- Simon MC, Keith B. 2008. The role of oxygen availability in embryonic development and stem cell function. *Nature Reviews Molecular Cell Biology* 9: 285-296. DOI: [10.1038/nrm2354](https://doi.org/10.1038/nrm2354)
- Stamati K, Mudera V, Cheema U. 2011. Evolution of oxygen utilization in multicellular organisms and implications for cell signalling in tissue engineering. *Journal of Tissue Engineering* 2: 204173141143236. DOI: [10.1177/2041731411432365](https://doi.org/10.1177/2041731411432365)
- Vermehren A, Langlais KK, Morton DB. 2006. Oxygen-sensitive guanylyl cyclases in insects and their potential roles in oxygen detection and in feeding behaviors. *Journal of Insect Physiology* 52: 340-348. DOI: [10.1016/j.jinsphys.2005.12.001](https://doi.org/10.1016/j.jinsphys.2005.12.001)
- Vermehren-Schmaedick A, Ainsley JA, Johnson WA, Davies SA, Morton DB. 2010. Behavioral Responses to Hypoxia in *Drosophila* Larvae Are Mediated by Atypical Soluble Guanylyl Cyclases. *Genetics* 186: 183-196. DOI: [10.1534/genetics.110.118166](https://doi.org/10.1534/genetics.110.118166)
- Yang CP, Fu CC, Sugino K, Liu Z, Ren Q, Liu LY, et al., Lee. 2015. Transcriptomes of lineage-specific *Drosophila* neuroblasts profiled via genetic targeting and robotic sorting. *Development* : 10.1242/dev.129163. DOI: [10.1242/dev.129163](https://doi.org/10.1242/dev.129163)

Funding:

DP was a post-doctoral researcher at the Instituto de Investigaciones Biológicas Clemente Estable, Ministerio de Educación y Cultura, Uruguay. Grants FVF-2019-05 (Dirección Nacional de Innovación, Ciencia y Tecnología) awarded to DP and FCE_1_2019_1_156160, awarded to RC and support from Agencia Nacional de Investigación e Innovación to DP and RC are acknowledged.

Author Contributions: Daniel Prieto: conceptualization, formal analysis, project administration, funding acquisition, writing - review editing, resources, visualization, investigation, writing - original draft. Boris Egger: conceptualization, methodology, supervision, writing - review editing. Rafael Cantera: conceptualization, funding acquisition, supervision, writing - review editing.

Reviewed By: Anonymous

Nomenclature Validated By: Anonymous

History: Received June 7, 2024 **Revision Received** July 29, 2024 **Accepted** August 2, 2024 **Published Online** August 11, 2024 **Indexed** August 25, 2024

Copyright: © 2024 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Prieto, D; Egger, B; Cantera, R (2024). Atypical soluble guanylyl cyclases control brain size in *Drosophila*. *microPublication Biology*. [10.17912/micropub.biology.001252](https://doi.org/10.17912/micropub.biology.001252)

