

The effect of GV-58, a calcium channel modifier, on synaptic transmission at the larval *Drosophila* and crayfish neuromuscular junctions

Jackson Schwamburger^{1*}, Kaitlyn Brock^{1*}, Robin Cooper^{1§*}

¹Biology, University of Kentucky, Lexington, Kentucky, United States

[§]To whom correspondence should be addressed: RLCOOP1@uky.edu

*These authors contributed equally.

Abstract

GV-58 is known to increase the opening time of the mammalian P-type calcium channel in presynaptic motor nerve terminals. GV-58 is suggested as a therapeutic agent for dampening the symptoms of amyotrophic lateral sclerosis. To further understand the mechanisms of GV-58 actions, the *Drosophila* and crayfish neuromuscular junctions were used as models. Their presynaptic calcium channels are a P-type based on pharmacology profiles. However, exposure of GV-58 (1mM) did not produce any consistent alteration in synaptic transmission in these two preparations. It is possible that the molecular structure of the P-type channels is different in the *Drosophila* and crayfish.




Figure 1. The effect of GV-58 on synaptic transmission at the *Drosophila* and crayfish neuromuscular junctions.:

(A) Dissection and recording of synaptic excitatory junction potentials (EJPs). (B) Four stimulation pulses every 10 seconds, the 1st and 4th EJP amplitudes were measured. (C) The amplitudes of the EJP saline and exposure to 0.1mM of GV-58 or to (D) 1 mM of GV-58. (E) The opener NMJ of the walking leg of a crayfish when stimulated at 40 Hz for 25 stimuli. (F) The 15th and 25th EJP amplitudes of a representative preparation in saline and (G) during exposure GV-58 (1mM).

Description

GV-58 is a compound that is suggested to target presynaptic voltage gated Ca^{2+} channels (VGCCs) of the “P-subtype”, or also known as Cav2.1. The VGCCs at the presynaptic motor nerve terminals of the larval *Drosophila* and crayfish neuromuscular junctions (NMJs) are both of a “P-subtype” (Araque et al., 1994; Kawasaki et al., 2004). GV-58 has been suggested to be used as a possible therapeutic agent to treat symptoms of amyotrophic lateral sclerosis (ALS). In the early onset of ALS synaptic efficacy is dampened. This is a critical time to help promote synaptic strength. Still there are no significant therapeutic treatments to retard the progression and prevention of motor neuron death. Selective enhancement of synaptic transmission at NMJs would aid in alleviating these early symptoms. Reports suggest that GV-58 is a modifier of the P-type calcium channels and prolongs the opening time of the channel when it opens due to voltage. Thus, more Ca^{2+} would enter the presynaptic nerve terminal and result in enhanced amount of transmitter release when the nerve is stimulated.

This investigation examined the effects of GV-58 on synaptic transmission at the model *Drosophila* and crayfish NMJs. Since synaptic transmission is graded, subtle changes in the presynaptic Ca^{2+} influx are readily measured by the amplitude and short-term facilitation (STF) of the excitatory postsynaptic potentials (EJPs) (Katz and Miledi, 1968;). STF is a transient increase in the amplitude of subsequent EJPs when more than one stimulus is rapidly given so that the residual Ca^{2+} will increase with each stimulus within the presynaptic terminal (Cooper and Cooper, 2009).

Exposure of GV-58 to the NMJs of the larval *Drosophila* body wall muscle and crayfish opener muscle did not reveal any consistent trend in altering synaptic transmission. The synaptic responses were mixed at the larval *Drosophila* NMJ, with some increasing the amplitude of the EJPs and some decreasing for both 0.1 mM and 1.0 mM exposure. However, for the crayfish NMJs there were no significant trends in the amplitudes of the EJPs within the trains for stimuli during the STF responses before or during exposure to GV-58. Since small changes in the extracellular Ca^{2+} concentration in the bathing saline as well as a buildup in the internal concentrations during STF are known to produce significant changes in the amplitudes of the EJPs in both preparations, it is concluded that GV-58 does not likely enhance the concentration of Ca^{2+} in the presynaptic terminals significantly during evoked stimulation to impact synaptic transmission.

GV-58 has been shown to have an enhancing effect in synaptic transmission at mouse NMJs (Li et al., 2023; Ojala et al., 2023), but it has also been shown to activate voltage gated Na^+ channels (Cho et al., 2022). Though the pharmacological profile indicates that the presynaptic calcium channels in the crayfish and *Drosophila* are deemed a P-type, it might not be relatable to the same “P-type” as classified in mammalian preparations. **A known blocker of P-type voltage-dependent Ca^{2+} channels is the venom of the funnel-web spider *Agelenopsis aperta* which contains the peptide omega-Aga-IVA.** This is why these channels were classified as such in the crayfish and other preparations (Araque et al., 1994). Also, physiological measures in the current voltage relationships of the crayfish motor neurons indicted a relationship between P-type calcium channels of mammals (Hong and Lnenicka, 1997; Tsien et al., 1988). The molecular identification of all the voltage gated ion channels within the presynaptic motor nerve terminals for the crayfish and *Drosophila* have yet to be fully identified and compared. Additionally, accessory proteins associated with the channels, which can be influenced by pharmacological agents, may also have a role in the function of the channels. Given the P-type Ca^{2+} channels of mammals are of the pharmacological nomenclature it would be of interest to obtain protein sequence comparisons among crayfish, *Drosophila* and mammalian models, along with accessory proteins, so further comparisons can be made as for potential screening of modulators for these channels to better understand the molecular mechanisms of action.

Methods

The experimental paradigm was to measure the effects before and during exposure to GV-58 on the amplitude and facilitation of the synaptic responses at the larval *Drosophila* and crayfish NMJs. GV-58 was dissolved in 0.1 % DMSO prior to adding to the saline. The saline control also contained 0.1 % DMSO.

Larval Drosophila melanogaster

Early 3rd instar Canton-S *Drosophila melanogaster* larvae were used for electrophysiological measures of muscle m6. Canton-S strain was obtained from Bloomington *Drosophila* Stock Center (stock # 64349). Transmembrane potentials were obtained with sharp intracellular electrodes (30 to 40 megaOhm resistance) filled with 3 M KCl. Fly saline haemolymph-like 3 (HL3) was used: (in mmol/L) 70 NaCl, 5 KCl, 20 MgCl_2 , 10 NaHCO_3 , 1 CaCl_2 , 5 trehalose, 115 sucrose, 25 N,N-bis(2-hydroxyethyl)-2-aminoethane sulfonic acid (BES) and pH at 7.1. An Axonclamp 2B (Molecular Devices, Sunnyvale, CA, USA) amplifier and 1 X LU head stage were used. The EJPs and spontaneous mEJPs were collected and analyzed with LabChart 7.0 (ADInstruments, USA).

Crayfish Opener muscle

Experiments were also performed using *Procambarus clarkii* bought directly from Kyle LeBlanc Crawfish Farms, Raceland, LA USA, 70394. Crayfish measuring 6–10 cm in body length and 12.5–25 g in body weight were used. Details of the dissection and electrophysiological recordings of the opener neuromuscular junction of the walking legs were previously described in video format (Cooper and Cooper, 2009). The crayfish saline used is in mM: 205 NaCl, 5.3 KCl, 13.5 CaCl₂·2H₂O, 2.45 MgCl₂·6H₂O, and 5 HEPES adjusted to pH 7.4. The excitatory axon innervating the opener muscle in the crayfish was stimulated in the meropodite by placing a branch of the leg nerve into a suction electrode connected to a Grass stimulator. Stimulation was with 40-Hz trains of 25 pulses in duration. Intracellular recordings were obtained in the same manner as described for the *Drosophila* preparation above. The nerve was stimulated at 40 Hz to facilitate the amplitudes of the EJPs to reliably measure them from baseline noise.

Acknowledgements: We thank the Bloomington *Drosophila* Stock Center for supplying the *Drosophila*.

References

- Araque A, Clarac F, Buño W. 1994. P-type Ca²⁺ channels mediate excitatory and inhibitory synaptic transmitter release in crayfish muscle.. Proceedings of the National Academy of Sciences 91: 4224-4228. DOI: [10.1073/pnas.91.10.4224](https://doi.org/10.1073/pnas.91.10.4224)
- Cho HY, Chen PC, Chuang TH, Yu MC, Wu SN. 2022. Activation of Voltage-Gated Na⁺ Current by GV-58, a Known Activator of CaV Channels. Biomedicines 10: 721. DOI: [10.3390/biomedicines10030721](https://doi.org/10.3390/biomedicines10030721)
- Cooper AS, Cooper RL. 2009. Historical View and Physiology Demonstration at the NMJ of the Crayfish Opener Muscle. Journal of Visualized Experiments : 10.3791/1595. DOI: [10.3791/1595](https://doi.org/10.3791/1595)
- Hong SJ, Lnenicka GA. 1997. Characterization of a P-Type Calcium Current in a Crayfish Motoneuron and Its Selective Modulation by Impulse Activity. Journal of Neurophysiology 77: 76-85. DOI: [10.1152/jn.1997.77.1.76](https://doi.org/10.1152/jn.1997.77.1.76)
- Katz B, Miledi R. 1968. The role of calcium in neuromuscular facilitation. The Journal of Physiology 195: 481-492. DOI: [10.1113/jphysiol.1968.sp008469](https://doi.org/10.1113/jphysiol.1968.sp008469)
- Kawasaki F, Zou B, Xu X, Ordway RW. 2004. Active Zone Localization of Presynaptic Calcium Channels Encoded by the *cacophony* Locus of *Drosophila*. The Journal of Neuroscience 24: 282-285. DOI: [10.1523/JNEUROSCI.3553-03.2004](https://doi.org/10.1523/JNEUROSCI.3553-03.2004)
- Li Y, Patel M, Baroudi J, Wu M, Gatti S, Liang M, et al., Meriney. 2023. A cross-sectional study of ageing at the mouse neuromuscular junction and effects of an experimental therapeutic approach for dynapenia. The Journal of Physiology 601: 4135-4150. DOI: [10.1113/JP284749](https://doi.org/10.1113/JP284749)
- Llinás R, Sugimori M, Lin JW, Cherksey B. 1989. Blocking and isolation of a calcium channel from neurons in mammals and cephalopods utilizing a toxin fraction (FTX) from funnel-web spider poison.. Proceedings of the National Academy of Sciences 86: 1689-1693. DOI: [10.1073/pnas.86.5.1689](https://doi.org/10.1073/pnas.86.5.1689)
- Ojala KS, Kaufhold CJ, Davey MR, Yang D, Liang M, Wipf P, Badawi Y, Meriney SD. 2023. Potentiation of neuromuscular transmission by a small molecule calcium channel gating modifier improves motor function in a severe spinal muscular atrophy mouse model. Human Molecular Genetics 32: 1901-1911. DOI: [10.1093/hmg/ddad019](https://doi.org/10.1093/hmg/ddad019)
- Tsien RW, Lipscombe D, Madison DV, Bley KR, Fox AP. 1988. Multiple types of neuronal calcium channels and their selective modulation. Trends in Neurosciences 11: 431-438. DOI: [10.1016/0166-2236\(88\)90194-4](https://doi.org/10.1016/0166-2236(88)90194-4)

Funding: We thank alumni of the research group for funding this project.

Author Contributions: Jackson Schwamburger: conceptualization, formal analysis, investigation, methodology, supervision, validation, visualization, writing - original draft, writing - review editing. Kaitlyn Brock : conceptualization, formal analysis, investigation, methodology, validation, visualization, writing - original draft, writing - review editing. Robin Cooper: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, software, supervision, validation, visualization, writing - original draft, writing - review editing.

Reviewed By: Ulises Ricoy

Nomenclature Validated By: Anonymous

History: Received October 28, 2024 **Revision Received** December 4, 2024 **Accepted** December 22, 2024 **Published Online** December 23, 2024 **Indexed** January 6, 2025

Copyright: © 2024 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any

12/23/2024 - Open Access

medium, provided the original author and source are credited.

Citation: Schwamburger, J; Brock , K; Cooper, R (2024). The effect of GV-58, a calcium channel modifier, on synaptic transmission at the larval *Drosophila* and crayfish neuromuscular junctions. microPublication Biology. [10.17912/micropub.biology.001402](https://doi.org/10.17912/micropub.biology.001402)